


Sarajevo

ICABB 2017 CONGRESS



**International Congress on
Advances in Bioscience and
Biotechnology**



BOOK OF ABSTRACTS

OCTOBER 25-29 2017

www.icabb.eu

1 Edited by

2 İlker Camkerten

3 Güzin Camkerten

4

5 Published, 2017

6 This work is subject to copyright. All rights are reserved, whether the whole
7 or part of the material is concerned. Nothing from this publication may be
8 translated, reproduced, stored in a computerized system or published in any
9 form or in any manner, including, but not limited to electronic, mechanical,
10 reprographic or photographic, without prior written permission from the
11 publisher.

12

13 www.icabb.eu

14 info@icabb.eu

15

16 The individual contributions in this publication and any liabilities arising
17 from them remain the responsibility of the authors.

18 The publisher is not responsible for possible damages, which could be a
19 result of content derived from this publication.

20

21

22 ISBN: (in progress)

23

24 24/11/2017

25

medipres
YAYINCILIK LTD. ŞTİ.

26

27

28 Dear Scientist,

29 The first International Congress on Advances in BioScience and
30 Biotechnology (icabb) was organized in Sarajevo / Bosnia and Herzogoniva.
31 We are very happy for organizing this congress in such a beautiful city and
32 country that we have strong historical ties.

33 We wanted to make this conference little bit special by bringing scientist
34 together from different disciplines of bioscience area and also to open new
35 research and cooperation fields for them. In this sense, we desired to bring
36 the distinguished scientist together to get know each other and to develop
37 and implement new joint projects.

38 The scientist joined the congress was from different country and mostly from
39 Turkey. The total number of submission were 103 and after a careful
40 evaluation 82 submissions were accepted by our scientific committee and 23
41 of them were accepted as poster presentation and 59 of them were accepted
42 as oral presentation and all those presentation was taken place in the
43 conference booklet.

44 We would like to send our special thanks to Mr. Musa Köse and Mr. İsmet
45 Uzun, ZENITH Group workers for their special efforts. And finally the most
46 importantly I would like to thank to all the participants individually who
47 came from far away to join this conference.

48

49

50 İlker Camkerten
51 Editor

52

53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78

ORGANIZATION COMMITTEE

Congress Chair

Assoc. Prof. Dr. İlker CAMKERTEN

Deputy Chairmans

Prof. Dr. Suat DİKEL, *Responsible for Ichthyology*

Prof. Dr. Kerem URAL, *Responsible for Veterinary Bioscience*

Prof. Dr. Erdoğan UZLU, *Responsible for Wildlife and Ecology*

Assoc. Prof. Dr. Yavuz Selim ÇAKMAK, *Responsible for Molecular Biology*

Assoc. Prof. Dr. Yaqoub ASHHAB, *Responsible for Bioinformatic*

Secretaries of Congress

Members of the Committee

Assoc. Prof. Dr. Deniz ALIÇ URAL

Asst. Prof. Dr. Güzin CAMKERTEN

İsmet UZUN, Zenith Group

Anes Bekric, Zenith Group

Alma LIGATA, Europe Congress

- 79 **SCIENTIFIC COMMITTEE**
- 80 **Khaled Abdou**, Professor at Environmental Toxicology, Vice Dean Faculty of Postgraduate
81 Studies for Advanced Sciences (PSAS), Faculty of Veterinary Medicine, Beni Suef University.
82 **EGYPT**
- 83 **Mohamed Ragab ABDEL GAWWAD**, Assoc. Prof. Dr. at Faculty of Engineering and Natural
84 Sciences, Biological Science and Bioengineering, International University of Sarajevo,
85 **BOSNIA&HERZOGOVINA**
- 86 **Zbigniew ADAMIAK**, Assoc. Prof. Dr. at Warmia-Mazury University, Olsztyn, **POLAND**
- 87 **Betul Akcesme**, Asst. Professor at Genetics and Bioengineering, International University of
88 Sarajevo, **BOSNIA&HERZOGOVINA**
- 89 **Faruk Berat Akçeşme**, Asst. Prof. Dr. at Genetics and Bioengineering, International
90 University of Sarajevo, **BOSNIA&HERZOGOVINA**
- 91 **İbrahim AKIN**, Asst. Prof. Dr. at Dept. of Surgery, Fac. Vet. Med., Adnan Menderes
92 University, **TÜRKİYE**
- 93 **Afsheen ARİF**, Asst. Professor at The Karachi Institute of Biotechnology and Genetic
94 Engineering (KIBGE) University of Karachi, **PAKISTAN**
- 95 **Mehmet AVCI**, Prof. Dr. at Animal Nutrition & Nutritional Diseases, Fac. Vet. Med, Harran
96 University, **TÜRKİYE**
- 97 **Mustafa Oguzhan CAGLAYAN**, Assoc. Prof. Dr. at Nanotechnology Engineering
98 Department, Cumhuriyet University, Sivas, **TURKEY**
- 99 **Irena CELESKA**, Assist. Prof. Dr. at Dept. of Pathophysiology, Fac. Vet. Med., Ss. Cyril and
100 Methodius University, **MACEDONIA**
- 101 **Çenker Çağrı CINGİ**, Assoc. Prof. Dr. at Dept. of Internal Medicine, Fac. Vet. Med.,
102 Afyonkocatepe University, **TÜRKİYE**
- 103 **Hakan ÇELEBİ**, Asst Professor at Department of Environmental Engineering, University of
104 Aksaray, **TURKIYE**
- 105 **Suat DİKEL**, Prof. Dr. at Dept. of Aquaculture, University of Çukurova, **TÜRKİYE**
- 106 **Mohamed EL HADİDİ**, Asst. Prof. at Bioinformatics - Head of the Bioinformatics Research
107 Group, Nile University, **EGYPT**
- 108 **Hesham Ali Metwally Ali EL-ENSHASY**, Professor at Bioprocess Engineering Dept. Faculty
109 of Chemical and Energy Engineering, Universiti Teknologi Malaysia (UTM), **MALAYSIA**
- 110 **Hasan ERDOĞAN**, Asst. Prof. Dr. at Dept. of Internal Medicine, Fac. Vet. Med., Adnan
111 Menderes University, **TÜRKİYE**
- 112 **Erdal EROL**, Assoc. Prof. Dr. at Dept. of Microbiology, University of Kentucky, **USA**
- 113 **Mehmet GÜLTEKİN**, Dr. at Dept. of Internal Medicine, Fac. Vet. Med., Adnan Menderes
114 University, **TÜRKİYE**
- 115 **Ali Evren HAYDARDEDEOĞLU**, Asst. Prof. Dr. at Dept. of Internal Medicine, Fac. Vet.
116 Med., Aksaray University, **TÜRKİYE**
- 117 **Mesut KARAHAN**, Asst. Prof. at Dept. of Bioengineering, Faculty of Engineering and
118 Natural Sciences, Üsküdar University, İstanbul, **TÜRKİYE**

- 119 **Muhammed KATICA**, Assoc. Prof. Dr. at Veterinary Clinical Pathology, Sarajevo University,
120 **BOSNIA&HERZOGOVINA**
- 121 **Obaid Yousuf KHAN**, Professor Dr., [Department of Genetics](#), [University of Karachi](#),
122 Karachi, **PAKISTAN**
- 123 **Koycho KOEV**, Asst. Prof. Dr. at Stara Zagora University, **BULGARIA**
- 124 **Serdar PAŞA**, Prof. Dr. at Dept. of Internal Medicine, Fac. Vet. Med., Adnan Menderes
125 University, **TÜRKIYE**
- 126 **Lazo PENDOVSKI**, Prof. Dr. at Ss. Cyril and Methodius University, Skopje, **MACEDONIA**
- 127 **Mohamad Roji bin Sarmidi**, Prof. Dr. Institute Bioproduct Development, Universiti
128 Teknologi Malaysia, 81310 UTM Johor Bahru, **MALAYSIA**
- 129 **Fahim SHALTOUT**, Professor of Meat Hygiene, Food Safety, Food Quality and Control
130 slaughterhouse trainer, Benha University, **EGYPT**
- 131 **Przemysław SOBIECH**, Assoc. Prof. Dr. at University of Warmia-Mazury, Olsztyn, **POLAND**
- 132 **Ilia TSHACEV**, Prof. Dr. at Stara zagora University, **BULGARIA**
- 133 **Shah Ali UL QADER**, Professor, Dr., Industrial Biotechnology, [Department of Biochemistry](#),
134 [University of Karachi](#), **PAKISTAN**
- 135 **Hikmet ÜN**, Assoc. Prof. Dr. at Dept. of Virology, Fac. Vet. Med., University of Aksaray,
136 **TÜRKIYE**
- 137 **Karolina WRZESNIEWSKA**, PhD, DVM at Dept. of Internal Medicine, Fac. Vet. Med.,
138 University of Lublin, **POLAND**
- 139 **Kaan YILANCIOĞLU**, Asst. Professor at Dept. of Bioengineering, Faculty of Engineering and
140 Natural Sciences, Üsküdar University, İstanbul, **TÜRKIYE**
- 141 **Katarzyna ŻARCZYŃSKA**, Assoc. Prof. Dr. at University of Warmia-Mazury, Olsztyn,
142 **POLAND**
- 143

PROGRAM OVERVIEW	
October 25, 2017	
	Airport transfers (Only for BLUE - GREEN - ORANGE PARTICIPANTS ID HOLDERS)
16:00-21:30	Sarajevo City Tour (Only for BLUE PACKAGE ID HOLDERS) MEETING IN THE HOTEL LOBBY
October 26, 2017	
08:00-09:30	Registration
09:30-09:45	Opening remarks - Welcome address - Opening speech
09:45-10:15	Plenary Session - Prof. Dr. Özcan Erel - Invited Speaker
10:15-10:30	Break Coffee &Tea
10:30-11:30	Session 1
11:30-11:50	Break Coffee &Tea
11:50-12:50	Session 2
13:00-14:00	LUNCH IN THE HOTEL RESTAURANT (Only for BLUE - GREEN - ORANGE Package ID Holders)
14:00-14:50	Session 3
14:50-15:10	Break Coffee &Tea Poster Session I
15:10-16:00	Session 4
16:00-16:20	Break Coffee &Tea Poster Session II
16:20-17:10	Session 5
17:10-17:30	Break Coffee &Tea Poster Session III
17:30-18:20	Session 6
19:30-20:30	Dinner in the Hotel Restaurant (Only for BLUE PACKAGE ID HOLDERS)
October 27, 2017	
08:00-09:00	Breakfast (Only for Blue and Green Package ID Holders)
09:00-09:50	Session 7
09:50-10:10	Break Coffee &Tea Poster Session IV
10:10-11:00	Session 8
11:00-11:20	Break Coffee &Tea Poster Session V
11:20-12:00	Session 9
12:00-13:30	Lunch (Only for BLUE - GREEN - ORANGE PARTICIPANTS ID HOLDERS)
13:30-14:20	Session 10
14:20-14:40	Break Coffee &Tea Poster Session VI

14:40-15:20	Session 11
15:20-15:40	Break Coffee & Tea Poster Session VII
15:40-16:10	Session 12
16:30-17:00	Closing Session: Closing remarks - Announcements - Closing speech
20:00-22:00	Gala Dinner (Only for BLUE - GREEN - ORANGE PARTICIPANTS ID HOLDERS) MEETING IN THE HOTEL LOBBY
October 28, 2017	
Social Cultural Tour to Mostar - Blagaj - Pocitelj (Only for BLUE PACKAGE ID HOLDERS)	
08:00	Breakfast (Only for Blue and Green Package ID Holders)
08:30	Departure from the Hotel
13:00	Lunch
19:00	Dinner
21:00	Departure to the Hotel
October 29, 2017	
08:30	Breakfast (Only for Blue and Green Package ID Holders)
09:00 - 12:00	Tunnel of Hope and Vrelo Bosna Tours
	Departures to Airport (Only for BLUE - GREEN - ORANGE PARTICIPANTS ID HOLDERS)

144

145

ORAL PRESENTATIONS OCTOBER 26, 2017

in honour of Prof. Özcan Erel

SESSION I

Chairmans: Prof. Dr. Özcan Erel - Prof. Mustafa Numan Bucak

10:30	20 - Overexpression of PHAC1 gene encoding PHA synthase 1 in Pseudomonas aureofaciens for overproduction of mcl-PHA. Emre Erden , Serap Evran, Nurdan Kasıkara Pazarlıoğlu.
10:40	26 - Association of MnSOD gene polymorphism and SOD activity and the risk of coronary artery disease in patients with non-diabetic metabolic syndrome. Seda Güleç Yılmaz , Atike Tekeli Kunt, Orhan Fındık, Selim İsbir, Turgay İsbir.
10:50	39 - The effect of ankaferd blood stopper and polyvinyl pyrrolidone on the model organism. Mehmet Okan Erdal , Eda Güneş, Lokman Gemi.
11:00	60 - D-lactic acid production from corn cob using L-LDH gene deficient Lactobacillus helveticus OZH12. Harun Önlü , Saadet Baltacı, Haluk Hamamcı, Özlem Osmanağaoğlu.
11:10	64 - Protective effect of omega-3 fatty acids on msg induced hippocampal neuronal apoptosis in the adolescent rats. Nurcan Umur , S. Gülşen Gürgen.
11:30	BREAK COFFEE/TEA

SESSION II

Chairman: Prof. Bülent Topcuoğlu

11:50	72 - Green synthesis of silver nanoparticles using Thymbra spicata var. spicata L. leaf extract and evaluation of their biological activities. Fatih Erci , Rabia Çakır Koç.
12:00	105 - The effects of prophylactic and therapeutic sulfasalazine in liver of experimental endotoxemic rats. Hatice Eser Fakı , Gonca Sönmez, Emre Bahçıvan, Burak Dik.
12:10	22 - The effects of silymarin on the cell viability in F-98 cell culture. Efe Kurtdede , Remzi Soner Cengiz, Ayşenur Gök, Ufuk Kaya, Görkem Kısmalı.
12:20	56 - Comparison of the beneficial effects of vitamin e and n-acethyl cysteine supplementations with therapeutic ultrasound application on the treatment of muscle injury of rats. Ayşe Merve Tat , Halil Ozkol, Yasın Tuluçe.
12:30	57 - Investigation of biochemical and histopathological effects of vitamin C, selenium and therapeutical ultrasound application on muscular injury formed in rats. Necati Muhammed Tat , Yasın Tuluçe, Mehmet Kara, Ayşe Merve Tat, Halil Ozkol.
12:40	54 - Effects of Metoclopramide in Maternal Rat Brain: Use to Increase of Breast Milk Production. Oya Sayın , Seren Gülşen Gürgen.

13:00	LUNCH IN THE HOTEL RESTAURANT (Only for BLUE - GREEN - ORANGE Package ID Holders)
SESSION III	
Chairman: Prof. Mehmet Kubilay Önal	
14:00	18 - Plant nutrient status of sour cherry (prunus cerasus l.) cultivars grown in Aegean region of Turkey. Bülent Topcuoğlu , M. Kubilay Onal.
14:10	19 - Characterization and evaluation of sour cherry (prunuscerasus l.) Genetic resources in Aegean region of Turkey. Mehmet Kubilay Önal , Bülent Topcuoğlu.
14:20	33 - The effect of spent mushroom compost on the growth, mineral nutrition and heavy metal status of lettuce (lactuca sativa l.) and spinach (spinaceae oleraceae l.). Bülent Topcuoğlu , M. Kubilay önal.
14:30	34 - The clonal selection breeding of akça pear (pyrus communis l.) in Aegean region of Turkey. Mehmet Kubilay Önal , Bülent Topcuoğlu.
14:40	85 - Shoot regeneration of Hypericum perforatum L. from axillary buds. Şeyma Önlü , Hussein Abdullah Ahmed Ahmed, Serkan Uranbey.
14:50	BREAK COFFEE / TEA Poster Session I (P1- P2- P5- P6)
SESSION IV	
Chairman: Assoc. Prof. Hidayet Argun	
15:10	21 - Modelling of phenazine-1-carboxylic acid production by pseudomonas chlororapsis subsp. aureofaciens in submerged cultures and investigation of its antimicrobial and anticancer activity. Kübra Aslan , Emre Erden.
15:20	27 - Hydrogen gas production using hot spring microflora. Ayşe Çalkaya, Handan Han, İkbal Ören, Nurçe Keskin, Zeynep Karaoğlan, Muhammed Enes Mıynat, İsmail Görgül, Hidayet Argun
15:30	24 - Effects of initial inoculum culture concentration on dark fermentative hydrogen gas production from waste paper towel hydrolysate. Hidayet Argun , Özgür Gökdemir, İsmail Görgül, Zeynep Saylam, Atakan Erdoğan, Gülizar Onaran, Hidayet Argun
15:40	28 - Investigation of biosorption kinetics of an agricultural residual on dye removal. Levent Gürel .
15:50	29 - Investigation of nickel biosorption by an agricultural residual. Levent Gürel , Tuğba Meşe, Arife Şeker, Aslı Kılıç, Ali Serkan Yıldırım.
16:00	BREAK COFFEE / TEA Poster Session II (P11- P20- P21)

SESSION V

Chairman: Asst. Prof.	
16:20	74 - A Deep convolutional neural network model for detecting diabetic retinopathy. Buse Melis Özyıldırım , Mustafa Oral, Serkan Kartal.
16:30	73 - Measuring nitrate consumption by <i>C. vulgaris</i> using NO ₃ - ion selective electrode (ISE). Vildan Erci , Didem Özçimen.
16:40	87 - Sulfasalazine may prevent disseminated intervacular coagulation in experimental endotoxemic rats. Emre Bahçivan , Hatice Eser Faki, Burak Dik.
16:50	30 - A new humanized mouse model for auto-immune myocarditis and its use to devise immunomodulation therapy. Mehmet Emrah Şelli , David C. Wraith, Andrew C. Newby.
17:00	31 - Autoimmunity to HSP60 during diet induced obesity in mice. Mehmet Emrah Şelli , David C. Wraith, Andrew C. Newby.
17:10	BREAK COFFEE / TEA Poster Session III (P4- P7- P8)

SESSION VI

Chairman: Asst. Prof. Numan Gözübenli	
17:30	107 - Pharmacokinetics of ceftriaxone following single ascending intravenous administration in sheep. Duygu Durna Çorum, Orhan Çorum, Hatice Eser Faki , Emre Bahçivan, Ayşe Er, Kamil Üney
17:40	106 - Shear bond strength of ceromer bonded to zirconium oxide infrastructure treated with different types of lasers. Oğuzhan Gorler, Ihsan Hubbezoglu, Melih Ulgey, Recai Zan, Serra Kutlu .
17:50	76 - Memory-enhancing, anxiolytic and antidepressant effects of inhaled <i>Achillea pseudoaleppica</i> essential oil on scopolamine-induced amnesic rats. Emel Akbaba , Eyüp Bağcı.
18:00	111 - The effects of <i>Juniperus oxycedrus</i> ssp <i>oxycedrus</i> essential oil on memory, anxiety and depression in amnesic rats. Emel Akbaba , Safeen Ahmed Othman, Eyup Bağcı.
18:10	108 - Doxorubicin loaded clamshell type four dithiol-o-carborane bridged metallophthalocyanines for photodynamic therapy. Berke Bilgenur Şener , Sevil Şener, Abdullah Tahir Bayraç.

in honour of Prof. Özcan Erel

SESSION VII

Chairman: Asst. Prof. Hasan Türe

09:00	8 - Antimicrobial Properties of Alginate and Zein Films Containing Natamycin against <i>Penicillium camemberti</i> and <i>Aspergillus niger</i> . Ömer Faruk Çelik , Gökçe Sarıtaş, Bekir Gökçen Mazı, Hasan Türe.
09:10	9 - Alginate beads for the removal of dyes from aqueous solution. Hasan Türe .
09:20	2 - Influence of trolox and alpha-lipoic acid on post-thawed pırlak ram sperm parameters and dna damage in non-breeding season - Fatih Avdatek , Deniz Yeni, Muhammed Kürşad Birdane, Mustafa Gündoğan.
09:30	13 - A cyclopia case with otocephaly in a pırlak lamb. Muhammed Kürşad Birdane , Oktay Yılmaz, Mehmet Uçar.
09:40	96 - Comparision of prostaglandin F ₂ α treatment in the Ovsynch protocol 6 or 7 days after first GnRH in non lactating Anatolian Buffaloes. Muhammed Kürşad Birdane , Fatih Avdatek, Hacı Ahmet ÇELİK.
09:50	BREAK COFFEE / TEA Poster Session IV (P3- P12- P15)

SESSION VIII

Chairman: Asst. Prof. Muhammed Kürşad Birdane

10:10	10 - mRNA expression of muscarinic receptors on human mesenchymal stem cells. Arash Alizadeh Yegani , Erkan Maytalan, İlknur Kozanoglu, Fazilet Aksu.
10:20	55 - The study of the effect on the bone tissue depending on the age gıngko biloba extract and growth factors. Seren Gülşen Gürgeın , Gülce Naz Yazıcı.
10:30	66 - A comparative investigation of the effects of various antiepileptic agents in the retina and corneal layers. Ayşe Tuç Yücel , Seren Gülşen Gürgeın, Ali Cansu.
10:40	91 - Ethanol induced changes in the expression of selected genes downstream to Nrf2 pathway in granulosa cells in vitro. Md. Mahmudul Hasan Sohel , Mustafa Abdulkareem Salman, Abdurrahman Ayvaz.
10:50	98 - Time and dose-dependent effects of oxidative stress induced by hydrogen peroxide in ovarian granulosa cells. Md. Mahmudul Hasan Sohel , Yusuf Konca, Mehmet Ulaş Çınar.
11:00	BREAK COFFEE / TEA Poster Session V (P9- P10- P13- P14)

SESSION IX

	Chairman: Prof. Nurettin Gülşen
11:20	44 - Effects of entodinium caudatum monoculture inoculation on in vitro fermentation, methane production and prevention of sub-acute ruminal acidosis. Nurettin Gülşen , Huzur Derya Arık, Mustafa Selçuk Alataş, Muhammad Naeem Tahir.
11:30	11 - Protective Effect of Colon Targeted Linalool and Lycopene Against Acetic Acid Induced Ulcerative Colitis in Rats. İbrahim Ozan Tekeli , Mustafa Yipel, Abdullah Aslan, Songül Çeribaşı, Ahmet Ateşşahin, Fatih Sakin
11:40	62 - Effect of feeding different protein levels on preference and some blood values in dogs. Fatma İnal, Oğuzhan Kahraman , Mustafa Selçuk Alataş, Şeref İnal, Mustafa Uludağ.
11:50	63 - Comparison of pelleted and extruded dog foods. Fatma İnal, Mustafa Selçuk Alataş, Oğuzhan Kahraman , Şeref İnal, Mustafa Uludağ.
12:00	LUNCH IN THE HOTEL RESTAURANT (Only for BLUE - GREEN - ORANGE Package ID Holders)
SESSION X	
	Chairman: Prof. Feride Koc
13:30	61 - Investigation of possibilities of using mobile bags in determining digestibility in dog foods. Mustafa Selçuk Alataş , Oğuzhan Kahraman, Fatma İnal, Nurettin Gülşen.
13:40	14 - Ameliorating effect of kisspeptin-10 on methotrexate-induced testicular oxidative stress and sperm quality in rats. Mehmet Güvenç , Mesut Aksakal.
13:50	92 - The effects of chrysine on cypermethrin-induced acute toxication in rainbow trout (<i>Oncorhynchus mykiss</i>). Özge Cerit, Feride Koç .
14:00	93 - The protective effects of Anethum graveolens on paracetamol-induced acute toxication in mice. Serap Korkmaz, Feride Koc
14:10	65 - The effect of heat treatment applied to cereals used in dog foods on gelatinization and digestibility of starch. Fatma İnal, Abdullah Özbilgin , Mustafa Selçuk Alataş, Oğuzhan Kahraman.
14:20	BREAK COFFEE / TEA Poster Session VI (P17- P 18- P19- P23)
SESSION XI	
	Chairman: Asst. Prof. Mehmet Güvenç
14:40	99 - Antimicrobial application of green synthesis of zinc oxide nanoparticles by euphorbia macroclada extract. Suzan Sahin Dogan , Aytac Kocabas.
14:50	70 - Some reproductive parameters in primiparous awassi sheep. Şeref İnal, Behiç Coşkun, Dursun Ali Dinç, Mustafa Çam .

15:00	71 - Pair housing for calves. Mustafa am , eref İnal.
15:10	78 - Effects of Thymol and Carvacrol on Sperm Quality and Antioxidant System. Mehmet Gven , Mustafa Cellat, İshak Gkek, İlker Yava, ule Yurdal zsoy.
15:20	BREAK COFFEE / TEA Poster Session VII (P16- P22)
SESSION XII	
Chairman: Asst. Prof. Gzin Camkerten	
15:40	109 - In vitro maturation in mouse. Mustafa Numan Bucak , Ali Erdem ztrk
15:50	89 - Investigation of iron, copper and zinc levels in sheep in district of Akakale. Metin Boz, İlker Camkerten .
16:00	88 - An investigation of blood iron, copper and zinc levels in sheep at region of Suru. İbrahim Halil elik, İlker Camkerten, Guzin Camkerten .
16:10	3 - Scalable fabrication of electrodes of tribo nanosensors based on highly uniform colloidal nano arrays. Numan Gzbenli . (Video Conference)

POSTER PRESENTATIONS OCTOBER 5 2017

1	110 - DNA library design for SELEX using bioinformatic tools. Berke Bilgenur Şener , Deniz Yiğit, Abdullah Tahir Bayraç.
2	25 - Analysis of protein and mRNA expressions of CYP1A1 and CYP2E1 enzymes in liver, colon and prostate cancer cell lines to study drug and carcinogen metabolism. Emre Evin , Serdar Karakurt, Orhan Adalı, Merve Akkulak.
3	35 - Comparative evaluation of small t gene and large t gene based real-time pcr methods for the detection of the bk and jc viruses in clinical samples. Uğur Tüzüner, Mehmet Özdemir , Murat Şevik, Mahmut Baykan.
4	41 - Effect of blood hormone levels and body condition score in success of artificial insemination in cow. İlker Yavaş, İshak Gökçek , Ece Koldaş, Tuğba Korkmaz Yavaş.
5	47 - The hemoglobinopathy prevalence of the patients with suspected hemoglobinopathy who have applied to Afyon Kocatepe University Practice and Research Hospital. Ahmet Kahraman , Mustafa Örmeci, Zeliha Çelik, Muhammed Kaya.
6	49 - Antioxidant properties of dietary flavonoids. Ahmet Kahraman .
7	52 - Investigation of İmmunological İnfertilit in non-fertilized cows. İlker Yavaş, Bestami Kemal Gümüşay, İshak Gökçek
8	59 - L-ldh gene knock-out of Lactobacillus helveticus OZH12. Harun Önlü , Marie-Clemence Duchene, Pascal Hols, Ozlem Osmanagaoglu.
9	68 - Effects of acetylsalicylic acid on inflammation and oxidative stress in rats with experimentally induced type 2 diabetes mellitus. Sefa Çelik , Serkan ŞEN.
10	69 - The effect of caffeic acid phenethyl ester (CAPE), a bioactive compound of propolis extract, on endoplasmic reticulum (ER) stress-induced apoptotic cell death in INS1-β cells. Sefa Çelik , Ahmet Kahraman, Ömer Hazman, Serkan Şen.
11	75 - Effect of hyperosmotic stress on MINPP1 expression. Fatih Erci , Mustafa Yöntem, Behiç Selman Erdoğan.
12	79 - Biochemical and haematological markers in patients with chronic HBV İnfections. Bilgen Özdemir , Ayşe Rüyeyda Uğur, Mehmet Özdemir, Mahmut Baykan.
13	82 - Investigation of parainfluenza virus caused lower respiratory tract infections in pediatric patients. Aysun Görkem, Ayşe Rüyeyda Uğur, Bahadır Feyzioğlu, Mehmet Özdemir, Mahmut Baykan .
14	83 - Investigation of parvovirus B19 IgM and IgG positivity rates in pediatric hematology patients. Aysun Görkem, Ayşe Rüyeyda Uğur, Fatma Taşbent, Mehmet Özdemir, Mahmut Baykan .
15	84 - Investigation of Bokavirus-related lower respiratory tract infections in pediatric patients. Ayşe Rüyeyda Uğur, Aysun Görkem, FATMA TAŞBENT, Bahadır Feyzioğlu, Mehmet Özdemir .

16	86 - An investigation of blood selenium, copper and zinc levels in sheep at region of Siverek. Mehmet Batmaz, Ilker Camkerten . Mehmet Batmaz, İlker Camkerten
17	97 - The post-hoc power analysis of forest productivity attributes in experimental study in central Bosnia. Azra Čabaravdić , Aida Ibrahimspahić, Mirsada Starčević.
18	101 - Rigorous identification of tissue specific genes in silico: revealing the interplay between cancer specific expression and tissue specific expression. Hatice Büşra Konuk , Alper Yılmaz.
19	102 - Generating lossless compression of genome scale k-mer frequency table as raster image. Hatice Büşra Konuk , Alper Yılmaz.
20	103 - Design and production of a lab-scale solar bioreactor. Vildan Erci , Didem Özçimen, İbrahim Işıldak.
21	122 - Evaluation of ischemia modified albumin levels in subclinical mastitis cows before and after the vaccination. Gaye Bulut, Güzin Camkerten , Özcan Erel.
22	104 - Diversity in mixed uneven aged forest of beech, fir and spruce on mountain Igman. Aida Ibrahimspahić, Azra Cabaravdic , Mirsada Starcevic.
23	Pair housing for calves. Mustafa Çam , Şeref İnal

149

150

interim

TABLE OF CONTENTS			
	Forewords		i
	Organization Committee		ii
	Program		iv
	Table of Contents		xvi
ORAL PRESENTATIONS			
ORAL 1	Influence of trolox and alpha-lipoic acid on post-thawed pirlak ram sperm parameters and dna damage in non-breeding season	Fatih Avdatek, Deniz Yeni, Muhammed Kürşad Birdane, Mustafa Gündoğan	2
ORAL 2	Scalable fabrication of electrodes of tribo nanosensors based on highly uniform colloidal nano arrays	Numan Gözübenli	3
ORAL 3	Alginate beads for the removal of dyes from aqueous solution	Hasan Türe	4
ORAL 4	Antimicrobial properties of alginate and zein films containing natamycin against penicillium camemberti and aspergillus niger	Ömer Faruk Çelik, Gökçe Sarıtaş, Bekir Gökçen Mazi, Hasan Türe	5
ORAL 5	Protective effect of colon targeted linalool and lycopene against acetic acid induced ulcerative colitis in rats	İbrahim Ozan Tekeli, Mustafa Yipel, Abdullah Aslan, Songül Çeribaşı, Ahmet Ateşşahin	6
ORAL 6	M-RNA expression of muscarinic receptors on human mesenchymal stem cells	Arash Alizadeh Yegani, Erkan Maytalan, İlknur Kozanoğlu, Fazilet Aksu	7
ORAL 7	Ameliorating effect of kisspeptin-10 on methotrexate-induced testicular oxidative stress and sperm quality in rats	Mehmet Güvenç, Mesut Aksakal	8
ORAL 8	Effects of thymol and carvacrol on sperm quality and antioxidant system	Mehmet Güvenç, Mustafa Cellat, İshak Gökçek, İlker Yavaş, Şule Yurdağül Özsoy	9
ORAL 9	A cyclopia case with otocephaly in a pirlak lamb	Muhammed Kürşad Birdane, Oktay Yılmaz, Mehmet Uçar	10
ORAL 10	Comparision of prostaglandin F ₂ α treatment in the Ovsynch protocol 6 or 7 days after first GnRH in non lactating Anatolian Buffaloes	Muhammed Kürşad Birdane, Fatih Avdatek, Hacı Ahmet Çelik	11
ORAL 11	Effect of feeding different protein levels on preference and some blood values in dogs	Fatma İnal, Oğuzhan Kahraman, Mustafa Selçuk Alataş, Şeref İnal, Mustafa Uludağ	12
ORAL 12	Comparison of pelleted and extruded dog foods	Fatma İnal, Mustafa Selçuk Alataş, Oğuzhan Kahraman, Şeref İnal, Mustafa Uludağ	13
ORAL 13	Overexpression of phaC1 gene encoding pha synthase 1 in pseudomonas aureofaciens for overproduction of mcl-pha	Emre Erden, Serap Evran, Nurdan Kaşıkara Pazarlıoğlu	14
ORAL 14	Modelling of phenazine-1-carboxylic acid production by pseudomonas chlororapsis subsp. aureofaciens in submerged cultures and investigation of its antimicrobial and anticancer activity	Kübra Aslan, Emre Erden	15
ORAL 15	The effects of silymarin on the cell viability in f-98 cell culture	Efe Kurtdede, Remzi Soner Cengiz, Ayşenur Gök, Ufuk Kaya, Görkem Kısmalı	16
ORAL 16	Effects of initial inoculum culture concentration on dark fermentative hydrogen gas production from waste paper towel hydrolysate	Hidayet Argun, Özgür Gökdemir, İsmail Görgül, Zeynep Sağlam, Gülizar Onaran	17
ORAL 17	Hydrogen gas production using hot spring microflora	Hidayet Argun, Ayşe Çalkaya, Handan Han, İkbal Ören, Zeynep Karaoğlan	18
ORAL 18	A new humanized mouse model for auto-immune myocarditis and its use to devise immunomodulation therapy	Mehmet Emrah Şelli, David C Wraith, Andrew C Newby	19
ORAL 19	Autoimmunity to hsp60 during diet induced obesity in mice	Mehmet Emrah Şelli, David C Wraith, Andrew C Newby	20
ORAL 20	Investigation of biosorption kinetics of an agricultural residual on dye removal	Levent Gürel	21
ORAL 21	Investigation of nickel biosorption by an agricultural residual	Levent Gürel, Tuğba Meşe, Arife Şeker, Aslı Kılıç, Alı Serkan Yıldırım	22
ORAL 22	Association of mnsod gene polymorphism and sod activity and the risk of coronary artery disease in patients with non-diabetic metabolic syndrome	Seda Güleç Yılmaz, Atike Tekeli Kunt, Orhan Fındık, Selim İşbir, Turgay İşbir	23
ORAL 23	Comparision of the beneficial effects of vitamin e and n-acetyl cysteine supplementations with therapeutic ultrasound application on the treatment of muscle injury of rats	Ayşe Merve, Halil Ozkol, Yasın Tuluce	24

ORAL 24	Investigation of biochemical and histopathological effects of vitamin c, selenium and therapeutical ultrasound application on muscular injury formed in rats	Necatı Muhammed Tat, Yasın Tuluçe, Mehmet Kara, Ayşe Merve Tat, Halil Ozkol	25
ORAL 25	The effect of ankaferd blood stopper and polyvinyl pyrrolidone on the model organism	Mehmet Okan Erdal, Eda Güneş, Lokman Gemi	26
ORAL 26	Investigation of parvovirus b19 Ig-m and Ig-g positivity rates in pediatric hematology patients	Aysun Görkem, Ayşe Rüveyda Uğur, Fatma Taşbent, Mehmet Özdemir, Mahmut Baykan	27
ORAL 27	Effect of blood hormone levels and body condition score in success of artificial insemination in cows	İlker Yavaşa, İshak Gökçek, Ece Koldaş, Tuğba Korkmaz Yavaş	28
ORAL 28	Effects of entodinium caudatum monoculture inoculation on in vitro fermentation, methane production and prevention of sub-acute ruminal acidosis	Nurettin Gülşen, Huzur Derya Arık, Mustafa Selçuk Alataş, Muhammad Naeem Tahir	29
ORAL 29	Effects of metoclopramide in maternal rat brain: use to increase of breast milk production	Oya Sayına, Seren Gülşen Gürgen	30
ORAL 30	The study of the effect on the bone tissue depending on the age ginkgo biloba extract and growth factors	Seren Gülşen Gürgen, Gülce Naz Yazıcı	31
ORAL 31	D-lactic acid production from corn cob using l-ldh gene deficient lactobacillus helveticus ozh12	Harun Önlü, Saadet Baltacı, Haluk Hamamcı, Ozlem Osmanağaoğlu	32
ORAL 32	Protective effect of omega-3 fatty acids on msg induced hippocampal neuronal apoptosis in the adolescent rats	Nurcan Umura, S. Gülşen Gürgen	33
ORAL 33	A deep convolutional neural network model for detecting diabetic retinopathy	Buse Melis Özyıldırım, Mustafa Oral, Serkan Kartal	34
ORAL 34	Shear bond strength of ceromer bonded to zirconium oxide infrastructure treated with different types of lasers	Oğuzhan Gorler, İhsan Hubbezoglu, Melih Ulgey, Recai Zan, Serra Kutlu	35
ORAL 35	Investigation of possibilities of using mobile bags in determining digestibility in dog foods	Mustafa Selçuk Alataş, Oğuzhan Kahraman, Fatma İnal, Nurettin Gülşen	36
ORAL 36	The effect of heat treatment applied to cereals used in dog foods on gelatinization and digestibility of starch	Fatma İnal, Abdullah Özbilgin, Mustafa Selçuk Alataş, Oğuzhan Kahraman	37
ORAL 37	A comparative investigation of the effects of various antiepileptic agents in the retina and corneal layers	Ayşe Tuç Yücel, Seren Gülşen Gürgen, Ali Cansu	38
ORAL 38	Sulfasalazine may prevent disseminated intervacular coagulation in experimental endotoxemic rats	Emre Bahçivan, Hatice Eser Fakı, Burak Dik	39
ORAL 39	Some reproductive parameters in primiparous awassi sheep	Şeref İnal, Behiç Coşkun, Dursun Ali Dinç, Mustafa Çam	40
ORAL 40	Green synthesis of silver nanoparticles using thymbra spicata var. Spicata l. Leaf extract and evaluation of their biological activities	Fatih Erci, Rabia Çakır Koç	41
ORAL 41	Measuring nitrate consumption by c. Vulgaris using no3- ion selective electrode (ise)	Vildan Erci, Didem Özçimen	42
ORAL 42	Memory-enhancing, anxiolytic and antidepressant effects of inhaled achillea pseudoaleppica essential oil on scopolamine-induced amnesic rats	Emel Akbaba, Eyüp Bağcı	43
ORAL 43	The effects of juniperus oxycedrus ssp oxycedrus essential oil on memory, anxiety and depression in amnesic rats	Emel Akbaba, Safeen Ahmed Othman, Eyüp Bağcı	44
ORAL 44	Shoot regeneration of hypericum perforatum l. From axillary buds	Şeyma Önlü, Hussein Abdullah Ahmed Ahmed, Serkan Uranbey	45
ORAL 45	Ethanol induced changes in the expression of selected genes downstream to nrf2 pathway in granulosa cells in vitro	Md. Mahmudul Hasan Sohel, Mustafa Abdulkareem Salman, Abdurrahman Ayvaz	46
ORAL 46	Time and dose-dependent effects of oxidative stress induced by hydrogen peroxide in ovarian granulosa cells	Md. Mahmudul Hasan Sohel, Yusuf Konca, Mehmet Ulaş Çınar	47
ORAL 47	The effects of chrysin on cypermethrin-induced acute toxication in rainbow trout (oncorhynchus mykiss)	Özge Cerit, Feride Koç	48
ORAL 48	The protective effects of anethum graveolens on paracetamol-induced acute toxication in mice	Serap Korkmaz, Feride Koç	49
ORAL 49	Antimicrobial application of green synthesis of zinc oxide nanoparticles by euphorbia macroclada extract	Suzan Şahin Doğan, Aytaç Kocabaş	50
ORAL 50	The effects of prophylactic and therapeutic sulfasalazine in liver of experimental endotoxemic rats	Hatice Eser Fakı, Gonca Sönmez, Emre Bahçivan, Burak Dik	51
ORAL 51	Pharmacokinetics of ceftriaxone following single ascending intravenous administration in sheep	Duygu Durna Çorum, Orhan Çorum, Hatice Eser Fakı, Emre Bahçivan, Ayşe Er, Kamil Üney	52
ORAL 52	Doxorubicin loaded clamshell type four dithiol-o-carborane bridged metallophthalocyanines for photodynamic therapy	Berke Bilgenur Şener, Sevil Şener, Abdullah Tahir Bayraç	53
ORAL 53	In vitro maturation in mouse	Mustafa Numan Bucak	54
ORAL 54	An investigation of blood iron, copper and zinc levels in sheep at region of Suruç	İbrahim Halil Çelik, İlker Camkerten, Güzin Camkerten, Tekin Şahin	55
ORAL 55	Investigation of iron, copper and zinc levels in sheep in district of Akçakale	Metin Boz, İlker Camkerten	56

POSTER PRESENTATIONS

POSTER1	DNA library design for selex using bioinformatic tools	Berke Bilgenur Şener, Deniz Yiğit,	58
---------	--------------------------------------------------------	------------------------------------	----

		Abdullah Tahir Bayraç	
POSTER 2	Analysis of protein and mrna expressions of cyp1a1 and cyp2e1 enzymes in liver, colon and prostate cancer cell lines to study drug and carcinogen metabolism	Emre Evin, Serdar Karakurt, Orhan Adalı, Merve Akkulak	59
POSTER 3	Comparative evaluation of small t gene and large t gene based real-time pcr methods for the detection of the bk and jc viruses in clinical samples	Uğur Tüzüner, Mehmet Özdemir, Murat Şevik, Mahmut Baykan	60
POSTER 4	Effect of blood hormone levels and body condition score in success of artificial insemination in cows	İlker Yavaşa, İshak Gökçek, Ece Koldaşç, Tuğba Korkmaz Yavaşd	61
POSTER 5	The hemoglobinopathy prevalence of the patients with suspected hemoglobinopathy who have applied to Afyon kocatepe university practice and research hospital	Ahmet Kahramana, Mustafa Örmeci, Zeliha Çelik, Muhammed Kaya	62
POSTER 6	Antioxidant properties of dietary flavonoids	Ahmet Kahraman	63
POSTER 7	Investigation of immunological infertility in non-fertilized cows	İlker Yavaşa, Bestami Kemal Gümüşay, İshak Gökçek	64
POSTER 8	L-ldh gene knock-out of lactobacillus helveticus ozh12	Harun Önlü, Marie-Clemence Duchene, Pascal Hols, Özlem Osmanağaoğlu	65
POSTER 9	Effects of acetylsalicylic acid on inflammation and oxidative stress in rats with experimentally induced type 2 diabetes mellitus	Sefa Çelik, Serkan Şen	66
POSTER 10	The effect of caffeic acid phenethyl ester (cape), a bioactive compound of propolis extract, on endoplasmic reticulum (er) stress-induced apoptotic cell death in ins1-β cells	Sefa Çelik, Ahmet Kahraman, Ömer Hazman, Serkan Şen	67
POSTER 11	Effect of hyperosmotic stress on minpp1 expression	Fatih Erci, Mustafa Yöntem, Behiç Selman Erdoğan	68
POSTER 12	Biochemical and haematological markers in patients with chronic hbv infections	Bilgen Özdemir, Ayşe Rüveyda Uğur, Mehmet Özdemir, Mahmut Baykan	69
POSTER 13	Investigation of parainfluenza virus caused lower respiratory tract infections in pediatric patients	Aysun Görkem, Ayşe Rüveyda Uğur, Bahadır Feyzioğlu, Mehmet Özdemir, Mahmut Baykan	70
POSTER 14	Investigation of parvovirus B19 IgM and IgG positivity rates in pediatric hematology patients	Aysun Görkem, Ayşe Rüveyda Uğur, Fatma Taşbent, Mehmet Özdemir, Mahmut Baykan	71
POSTER 15	Investigation of bokavirus-related lower respiratory tract infections in pediatric patients	Ayşe Rüveyda Uğura, Aysun Görkem, Fatma Taşbent, Bahadır Feyzioğlu, Mehmet Özdemir	72
POSTER 16	An investigation of blood selenium, copper and zinc levels in sheep at region of Siverek	Mehmet Batmaz, İlker Camkerten	73
POSTER 17	The post-hoc power analysis of forest productivity attributes in experimental study in central Bosnia	Azra Čabaravdić, Aida Ibrahimspahić, Mirsada Starčević	74
POSTER 18	Rigorous identification of tissue specific genes in silico: revealing the interplay between cancer specific expression and tissue specific expression	Hatice Büşra Konuk, Alper Yılmaz	75
POSTER 19	Generating lossless compression of genome scale k-mer frequency table as raster image	Hatice Büşra Konuk, Alper Yılmaz	76
POSTER 20	Design and production of a lab-scale solar bioreactor	Vildan Erci, Didem Özçimen, İbrahim Işıldak	77
POSTER 21	Evaluation of ischemia modified albumin levels in subclinical mastitis cows before and after the vaccination	Gaye Bulut, Güzin Camkerten, Özcan Erel	78
POSTER 22	Diversity in mixed uneven aged forest of beech, fir and spruce on mountain Igman	Ibrahimspahić, Azra Cabaravdic, Mirsada Starcevic	79
POSTER 23	Pair housing for calves	Mustafa Çam, Şeref İnal	80
FULL TEXT (FT) PRESENTATIONS			
FT 1 ORAL 56	The effect of spent mushroom compost on the growth, mineral nutrition and heavy metal status of lettuce (<i>lactuca sativa</i> l.) and spinach (<i>spinaceae oleraceae</i> l.)	Bülent Topcuoğlu, Mehmet Kubilay Önal	82
FT 2 ORAL 57	Plant nutrient status of sour cherry (<i>prunus cerasus</i> l.) Cultivars grown in aegean region of Turkey	Bülent Topcuoğlu, Mehmet Kubilay Önal	91
FT 3 ORAL 58	Characterization and Evaluation of Sour Cherry (<i>Prunus cerasus</i> l.) Genetic Resources in Aegean Region of Turkey	Mehmet Kubilay Önal, Bülent Topcuoğlu	98
FT 4 ORAL 59	The Clonal Selection Breeding of Akça Pear (<i>Pyrus communis</i> l.) in Aegean Region of Turkey	Mehmet Kubilay Önal, Bülent Topcuoğlu	108

153

154

155

156

157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184

interim

ORAL PRESENTATIONS

Oral - 1

185 **INFLUENCE OF TROLOX AND ALPHA-LIPOIC ACID ON POST-**
186 **THAWED PIRLAK RAM SPERM PARAMETERS AND DNA DAMAGE IN**
187 **NON-BREEDING SEASON**

188 **FATİH AVDATEK^a, DENİZ YENİ^a, MUHAMMED KÜRŞAD BİRDANE^a**
189 **MUSTAFA GÜNDOĞAN^a**

190
191 ^a Afyonkocatepe university

192
193 favdatek@aku.edu.tr

194 **The present study** was conducted to examine the protective roles of trolox and
195 alpha-lipoic on post-thawed Pırlak ram sperm parameters and DNA damage in non-
196 breeding season. Semen samples from 10 healthy Pırlak rams (2-3 years of age)
197 were used in the study. Six ejaculates for each rams were collected and pooled.
198 Pooled ejaculate, splitted into three equal aliquots and then diluted with Tris-based
199 extender at 37 °C containing trolox 1 mM, alpha-lipoic acid (ALA) 1 mM and no
200 antioxidant (control) were cooled to 5 °C and then frozen in 0.25 ml French straws.
201 Frozen straws were then thawed at 37°C for 20s in a water bath for evaluation.
202 **Sperm motility** was increased, acrosome rate and DNA damage were decreased
203 significantly ($p<0.05$) in ALA, head abnormal sperm rate and membrane integrity
204 were increased, acrosome rate and DNA damage were decreased significantly ($P <$
205 0.05) in trolox when compared to the control group. **Results** of this study suggest
206 that alpha-lipoic acid and trolox improve sperm parameters and DNA damage in
207 non-breeding season.

208

209 **Keywords:** Alpha-Lipoic Acid, Non-Breeding Season, Ram Sperm, Trolox

210

211

Oral - 2

212 **SCALABLE FABRICATION OF ELECTRODES OF TRIBO**
213 **NANOSENSORS BASED ON HIGHLY UNIFORM COLLOIDAL NANO**
214 **ARRAYS**

215

NUMAN GÖZÜBENLİ

216

Harran university

217

218

219

gnuman@harran.edu.tr

220 Self-assembled two-dimensional (2-D) nanoarrays have been extensively utilized for
221 fabricating a wide spectrum of functional periodic nanostructures for important
222 technological applications. Here we report a simple and scalable colloidal
223 transferring technique for fabricating highly ordered 2-D colloidal nanoarrays with
224 unusual non-close-packed structures on a large variety of substrates (e.g., plastics
225 and glass electrodes of triboelectric generators (TENG's)). A copolymer made of
226 mixture of 1:1 ratio of ethoxylated trimethylolpropane triacrylate (ETPTA) and
227 urethane (PU) monomers is used as the particle transferring agent to transfer
228 colloidal particles from a self-assembled 3-D colloidal crystals in a layer-by-layer
229 manner. **The unique combination** of optical transparency, flexibility, and
230 durability of the copolymer enables wafer-scale transfer of colloidal monolayers
231 onto both flat and curved surfaces, which are not easily available by current top-
232 down and bottom-up approaches for fabrication of electrodes of TENG's. **We also**
233 **show** that this method provides large-area production of hexagonally ordered
234 nanostructures (e.g., periodic nanoarrays for potential applications of TENG's)
235 which have important applications ranging from self-powered portable devices
236 harvesting electricity to biosensors.

237

238 **Keywords:** Triboelectric Nanosystems, Contact Electrification, Layer By Layer
239 Transfer Tech, Tribo Nanosensors

240

241

Oral - 3

242

ALGINATE BEADS FOR THE REMOVAL OF DYES FROM AQUEOUS SOLUTION

243

244

HASAN TÜRE

245

Ordu university

246

247

248

hasanture@odu.edu.tr

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

Several types of industries including paper, textile and rubber use significant amount of dye to color their products. However, contamination of water by dyes discharged from these industries is a significant threat to human and ecosystems since they are toxic and pose carcinogenic and mutagenic effects. Several types of technologies are available to remove dyes from water such as coagulation-flocculation, membrane filtration, ion exchange, and electrochemical techniques. On the other hand, most of these methods are expensive and can produce other waste products. Thus, there is a need to develop effective and environmentally friendly technologies to remove the dyes from water. **Adsorption** is one of the most preferred methods to remove organic and inorganic pollutants from wastewater owing to its simplicity of design, flexibility and ease of operation. Various low cost and easily available adsorbents have been used to remove dyes from water. **In this study**, alginate beads were prepared by ionic gelation method and the adsorption of methylene blue dye (MB) on wet and dry alginate beads was tested as a function of pH, contact time, initial MB concentration, and temperature in batch system. In addition, montmorillonite was also incorporated into the alginate beads and their characterization was performed. **Results** indicated that the amount of dye adsorbed is highly depended on the initial dye concentration, pH and temperature. Dry alginate beads showed higher adsorption capacity than wet beads. SEM-EDX analysis revealed that montmorillonite was existed as agglomerates. Obtained alginate beads could be used not only for removal dyes but also used to eliminate toxic metal ions from water.

Keywords: Alginate, Dye, Beads, Adsorption

274

275 **ANTIMICROBIAL PROPERTIES OF ALGINATE AND ZEIN FILMS**
276 **CONTAINING NATAMYCIN AGAINST PENICILLIUM CAMEMBERTI**
277 **AND ASPERGILLUS NIGER**

278 **OMER FARUK CELIK^a, GÖKCE SARITAŞ^b, BEKİR GÖKÇEN MAZİ^c, HASAN TÜRE^d**

279 **aaaaaaaaaaaaaaaaaaaaaaaaaaaaa**

280

281 ofarukcelik@gmail.com

282 **The decline in food sources** as a consequence of the increasing world population
283 and environmental pollution has made conservation of natural and processed foods
284 even more important. New packaging technologies as well as packaging and storage
285 techniques that provide longer shelf life should be developed. Edible films and
286 coatings obtained from renewable resources could be considered as promising
287 packaging materials since they are biodegradable, inexpensive and environmentally
288 friendly.

289 Kashar cheese is a cooked-curd cheese and it is usually classified in the same
290 group with Mozzarella cheese. One of the main issues concerning the quality and
291 shelf life of kashar cheese is mould spoilage. A recent methodology proposed to
292 maintain the food safety and to extend the shelf life of foods is incorporation of
293 natural antimicrobial agents into the biobased packaging materials. Natamycin
294 (NA) is a natural antifungal agent and has been approved as Generally Recognized
295 as Safe (GRAS) in many countries for surface application of cheese.

296 The goal of this study is to investigate the antifungal activities of alginate and zein
297 films containing different amounts of NA (0-2-5-10-20-40 mg) against *Penicillium*
298 *camemberti* and *Aspergillus niger* inoculated on the surface of kashar cheese
299 during refrigerated storage. Both zein and alginate films containing 20 and 40 mg
300 NA were found to be effective against *A. niger* with reductions of 1.41 and 1.5 logs
301 for zein and 0.65 and 1.21 logs for alginate. Zein films with 10, 20 and 40 mg NA
302 were found to decrease *P. camemberti* population by 1.69; 1.78 and 1.85 logs,
303 respectively. Conversely, alginate films incorporated with NA had weak inhibition
304 effect against *P. camemberti* during storage and no effect at the end.

305 Although both films incorporated with NA appear to be successful systems, zein
306 films could practically serve better to overcome the surface contamination of the
307 dairy products.

308

309 **Keywords:** Alginate, Zein, Edible Film, Cheese, Natmaycin

310

311
312 **PROTECTIVE EFFECT OF COLON TARGETED LINALOOL AND**
313 **LYCOPENE AGAINST ACETIC ACID INDUCED ULCERATIVE COLITIS**
314 **IN RATS**

315 **İBRAHİM OZAN TEKELİ^a, MUSTAFA YİPEL^b, ABDULLAH ASLAN^c**
316 **SONGÜL ÇERİBAŞI^d, AHMET ATEŞŞAHİN^e**

317
318 ^aMustafa Kemal uni., faculty of veterinary medicine, dept. of pharmacology and toxicology

319 ^bNamik Kemal uni., faculty of veterinary medicine, dept. of pharmacology and toxicology

320 ^cFirat university, faculty of science, department of biology

321 ^dFirat university, faculty of veterinary medicine, department of pathology

322 ^eFirat university, faculty of veterinary medicine, department of pharmacology and toxicology

323
324 ozantekeli@hotmail.com

325
326 The study was carried out with the aim of the investigation of protective effect of
327 conventional and colon targeted linalool and lycopene comparatively in rats with
328 experimental ulcerative colitis (UC). A total of 49 Wistar Albino rats (7 animals in
329 each group) were used. Conventional and colon targeted linalool (200 mg/kg) and
330 lycopene (10 mg/kg) were given to them for 7 days and then UC was induced using
331 4% acetic acid intrarectally. To exhibit the oxidative damage and antioxidant
332 activity and the inflammation status of the colon tissues that were taken at end of
333 the experimental model. Malondialdehyde (MDA), reduced glutathione (GSH),
334 catalase (CAT), glutathione peroxidase (GPx), tumor necrosis factor α (TNF- α),
335 interleukin 1 β (IL-1 β), interleukin 6 (IL-6) levels and cyclooxygenase 2 (Cox-2),
336 nuclear factor erythroid 2-related factor 2 (Nrf-2) and nuclear factor kappa B (NF-
337 kB) expression levels were examined. In addition, histopathological scoring was
338 performed in terms of depth and width of ulcerative lesions. Consequently, the
339 usage of targeted agents was found to be more effective than conventional usage
340 against to histopathologic, oxidative and inflammatory damage occurred in the
341 colon in the experimental UC model. Also, targeted lycopene was found to be more
342 effective than targeted linalool against acetic acid induced UC in rats.

343
344 **Keywords:** Colitis, Acetic Acid, Rat, Colon Targeting, Protective Effect

345

346

Oral - 6

347

MRNA EXPRESSION OF MUSCARINIC RECEPTORS ON HUMAN MESENCHYMAL STEM CELLS

348

349

ARASH ALIZADEH YEGANI^a, ERKAN MAYTALMAN^b, İLKNUR KOZANOGLU^c,
FAZİLET AKSU^d

350

351

^aMustafa Kemal uni., dept. of pharmacology and toxicology, faculty of veterinary medicine

352

^bAlanya Alaaddin Keykubat university, department of medical pharmacology

353

^cBaşkent university faculty of medicine, adult bone marrow transplant center

354

^dÇukurova university faculty of medicine department of medical pharmacology

355

356

357

arash.alizadeh82@gmail.com

358

Cells carry out their functional activities by communicating with each other through endogenous substances and receptors. For regeneration purposes, it may be possible to demonstrate the functional activities required for transplantation of new stem cells, where stem cells are placed by transporting appropriate receptors on cell surfaces. Demonstration of which receptors are present in mesenchymal stem cells will allow these cells to be used more appropriately for regeneration purposes. **In this study** the mRNA expression levels of M1, M2, M3, M4, M5 subtypes of muscarinic receptors in the 1st, 2nd and 3rd passages of Mesenchymal stem cells (MSCs) obtained from human fetal membrane (FM) and bone marrow (BM) were shown by RT-qPCR. In our study, a generally significant increase in FM groups was detected in M1 mRNA expression compared to the control group. Differently in the BM groups was seen as a decrease. Results were obtained significant decrease in the M5 mRNA expression the of BM groups compared to the control group. **These results** indicate that the cells obtained from different sources in that the expressions of muscarinic receptors behave differently and there are also some changes derived from the same sources cells according to the passages and differentiations.

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

Keywords: Mesenchymal Stem Cells, Muscarinic Receptors, RT-Qpcr

377

379 **AMELIORATING EFFECT OF KISSPEPTIN-10 ON METHOTREXATE-**
380 **INDUCED TESTICULAR OXIDATIVE STRESS AND SPERM QUALITY IN**
381 **RATS**

382 **MEHMET GÜVENÇ^a, MESUT AKSAKAL^b**

383

384 ^aMustafa Kemal university, veterinary science, department of physiology

385 ^bFırat university, veterinary science, department of physiology

386

387

vetguvenc@gmail.com

388 **The purpose** of this study is to determine effects of methotrexate (MTX) on
389 oxidative stress, antioxidative system and spermatogenesis and changes caused by
390 Kisspeptin-10 (Kiss) administration on these effects. **For this purpose**, total of 32
391 wistar albino rats were divided into groups as; group I control (n:6, saline), group II
392 MTX (n:10, i.p MTX 20 mg/kg single dose), group III (n:6, Kisspeptin-10, 50
393 nmol/kg, 10 days/dose), and group IV (n:10, MTX 20 mg/kg single dose +
394 Kisspeptin-10, 50 nmol/kg, 10 days/dose). **Spermatological parameters**
395 (including motility, epididymal sperm concentrations live-dead sperm rate and
396 morphological sperm abnormalities), biochemical parameters (MDA, GSH, GSH.Px
397 and CAT) of testicular tissue were evaluated. **At the end of the study**, MDA levels
398 increased by MTX applications in testicular tissue ($p < 0.05$) and showed significant
399 decrease in group IV by Kisspeptin administration ($p < 0.05$). The levels of GSH in
400 the testicular tissue were significantly increased in group IV ($p < 0.05$). While
401 testicular tissue GSH.Px levels decreased with MTX administration ($p < 0.05$), in
402 group IV was increased by Kisspeptin administration ($p < 0.05$). In spermatological
403 parameters, MTX administration (group II), decreased motility ($p < 0.001$), and
404 increased rates of abnormal spermatozoon ($p < 0.05$), in treatment group (group IV)
405 an improvement was observed with increased motility ($p < 0.001$), and decreased
406 abnormal spermatozoon ($p < 0.05$). **As a result**, MTX administration caused damage
407 on some spermatological parameters, increased oxidative stress when compared to
408 the control group. However, Kiss treatment mitigates these side effects when
409 compared to the MTX group. **It is concluded that** Kiss treatment may reduce
410 MTX-induced reproductive toxicity as a potential antioxidant compound.

411 **Keywords:** Kisspeptin, Methotrexate, Oxidative Stress, Sperm Quality

412

EFFECTS OF THYMOL AND CARVACROL ON SPERM QUALITY AND ANTIOXIDANT SYSTEM

MEHMET GÜVENÇ^a, MUSTAFA CELLAT^b, İSHAK GÖKÇEK^c, İLKER YAVAŞ^d
ŞULE YURDAGÜL ÖZSOY^e

^aMustafa kemal university, faculty of veterinary medicine

vetguvenc@gmail.com

In this study, the effects of different doses of thymol and carvacrol on sperm quality and antioxidant status were investigated. **For this purpose**, 49 rats were divided to 7 animals in each group; Control(saline), T-10; (Thymol 10 mg/kg), T-20; (Thymol 20 mg/kg), C-10; (Carvacrol 10 mg/kg), C-20; (Carvacrol 20 mg/kg), T+C-10; (Thymol 10 mg/kg + Carvacrol 10 mg/kg) and T+C-20; (Thymol 20 mg/kg + Carvacrol 20 mg/kg). The application lasted 10 weeks for all animals. **In the study**, spermatological parameters (including motility, concentration, morphological sperm abnormalities, live-dead sperm rate), biochemical parameters (MDA, GSH, GSH.Px, Catalase) and histopathological analyzes were evaluated. **As a result of the study**, MDA levels decreased compared to the control group in the group T-10, group T-20, group C-10 and group C-20 ($p < 0.001$). GSH levels were increased in group T-10 and in group T-20 compared to the control group ($p < 0,01$). GSH.Px levels were higher in group C-10 and in group C-20 than in the control group ($p < 0,05$). Catalase levels were increased in Group C-10 and Group C-20 compared to the control group ($p < 0,05$). **In spermatological parameters**, both spermatozoon density and motility were significantly higher in the group T-10, group T-20, group C-10, group C-20 and group T+C-10 compared to the control group ($p < 0,01$). There was a decrease in group T+C-20 in spermatozoon concentration, compared to control group ($p < 0,001$). In the live-dead sperm rate and abnormal spermatozoon ratio there was significantly decrease in group T-10, group T-20, group C-10, group C-20 compared with the control group ($p < 0.001$). **Histopathological analysis** of animals from testicular tissues revealed no differences among the groups. **In conclusion**, it was found that thymol and carvacrol treatment decreased oxidative damage, increased antioxidant status and showed improvement in spermatological parameters, while the combined use of these two active substances showed no improvement effect on the parameters mentioned.

Keywords: Antioxidant, Sperm Quality, Thymol, Carvacrol

Acknowledgements: This study is supported by Scientific Activities Support Program of Mustafa Kemal University

A CYCLOPIA CASE WITH OTOCEPHALY IN A PIRLAK LAMB

MUHAMMED KÜRŞAD BIRDANE, OKTAY YILMAZ, MEHMET UÇAR

Afyonkocatepe university

mkbirdane@yahoo.com

In this report, a case of cyclopia with otocephaly in a Pirlak lamb was presented which has not been reported previously in ruminants. A two years old-nulliparous Pirlak ewe was referred to the clinics of Afyon Kocatepe University, Faculty of Veterinary Medicine with the complaint of dystocia. Cesarean section was performed and a dead female lamb was extracted out from the uterus. The number and size of placentomes were reduced at the placental sites. Morphologic and macroscopic examination of the lamb demonstrated that there were some congenital anomalies, especially an eye (cyclopia) and otocephaly, at the head of lamb, however no other anomalies were defined at the other parts of body. On the other hand there were no whole mouth space, maxillar and mandibular bones, teeth (dysgnathia), no nose (arhinia) and tongue (aglossia). Nevertheless, there was a small hole at the bottom of the one eye opening to pharynx, oesophagus and trachea (microstomia). The left and right auricles were adherent together on the both sides of the small hole (synotia). **At necropsy**, the brain was not normal, rather small and edematous, having any brain lobes and gyrus. Examination of thoracic, abdominal and pelvic spaces revealed no abnormality

Keywords: Congenital Anomalies, Pirlak Lamb, Cyclopia, Otocephaly

480 **COMPARISON OF PROSTAGLANDIN F₂A TREATMENT IN THE**
481 **OVSYNCH PROTOCOL 6 OR 7 DAYS AFTER FIRST GNRH IN NON**
482 **LACTATING ANATOLIAN BUFFALOES**

483 **MUHAMMED KÜRŞAD BIRDANE^a, FATİH AVDATEK^b, HACI AHMET ÇELİK^c**

484

485 ^a Obstetrics and gynaecology, veterinary medicine, Afyonkocatepe university

486 ^b Reproduction and artificial insemination, veterinary medicine, Afyonkocatepe university

487 ^c Obstetrics and gynaecology, faculty of veterinary medicine, Afyonkocatepe university

488

489

mkbirdane@yahoo.com

490 **In this study**, Ovsynch versus modified Ovsynch protocols on follicle size at
491 insemination day and pregnancy rate. The total of 26 non lactating buffalo cows
492 whose ages were between 6-8 year old age, belonging to a private buffalo farm in
493 Afyonkarahisar were used in this study. In February, estrous synchronization of
494 Group 1 (n = 14) was performed with GnRH at day 0, PGF₂α at day 7, and GnRH at
495 day 9 (12 µg buserelin, Receptal®, 3 ml i.m). Estrous synchronization of Group 2
496 (n=11) was similar with Group 1 except PGF₂α treatment at day 6. The rate of
497 response to synchronization was determined at the time of insemination and the
498 pregnancy rates were determined at day 42 using transrectal ultrasonography.
499 Artificial insemination was performed using frozen thawed Italian buffalo semen,
500 16-20 h later last GnRH injection in Group 1 and 2. Ovulation investigation was
501 made after 1 day of insemination. The ovulatory follicle diameter (>1cm) were
502 found 1.30 ± 0.10 cm in Group 1 and 1.26 ± 0,12 cm in Group 2. The response to
503 Ovsynch in the groups were 78,5% (11/14) in Group 1 and 81,8 % (9/11) in Group
504 2. Pregnancy rates were found % 36,3 (4/11) in Group 1 and % 33,3 (3/9) in
505 Group 2. **As a result**, It is thought that there is a need for a larger number of
506 animals in groups to establish the difference between the day 6 and 7 PGF₂α
507 injection in Ovsynch protocols.

508

509 **Keywords:** Anatolian Buffaloes, Non Lactating, Ovsynch, Pregnancy Rate

510

511

Oral - 11

512

EFFECT OF FEEDING DIFFERENT PROTEIN LEVELS ON PREFERENCE AND SOME BLOOD VALUES IN DOGS

513

514

FATMA INAL^a, OGUZHAN KAHRAMAN^a, MUSTAFA SELCUK ALATAS^a

515

SEREF INAL^a, MUSTAFA ULUDAG^b

516

517

^aSelcuk university faculty of veterinary medicine, department of animal nutrition

518

^bBil-yem gida san. ve ltd. sti. Ankara, TURKEY

519

520

vetoguzhan90@gmail.com

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

This study was conducted to observe the effect of feeding different protein levels in dogs. The mix breed castrated young mature male dogs at approximately 15-30 kg live weights have been used. 4 extruded foods containing 18%, 21, 25 and 28 crude protein in dry matter were prepared in Bil-Yem facilities. **Foods** were eaten for a month. The dogs were divided into 4 similar groups, each consisting of 7 dogs according to locations in the unit, live weights and body condition scores. The animals were housed in individual compartments with a closed and a walking area. The foods were given at the rate of 1.8 times of maintenance once a day. The blood samples were taken at the beginning and end of the study. It was determined that BUN values were affected by protein level (P=0.002) and creatinine, total protein, albumin and phosphorus levels not. **Body** conditions were not affected by protein levels. **Thirty dogs** were used in two-pan palatability test. Two foods with 21% and 28% protein were tested according to dog's preference. Dogs have preferred the food with high protein at the rate of 67%. **As the** protein increases, the cost of the food increased.

536

537

Keywords: Dog Food, Protein, BUN, Preference

538

COMPARISON OF PELLETTED AND EXTRUDED DOG FOODS

FATMA INAL^a, MUSTAFA SELÇUK ALATAS^a, OGUZHAN KAHRAMAN^a, SEREF

INAL^a, MUSTAFA ULUDAG^b

^aSelçuk university faculty of veterinary medicine, department of animal nutrition

^bBil-yem gıda san. ve ltd. sti. Ankara, TURKEY

vetoguzhan90@gmail.com

In the study, 30 young adult mixed breed neutered male dogs weighing 15-30 kg were used. **One** of the two formulations, which had the same composition, was produced in the form of a pelleted and the other an extruded in a private factory. Feeding experiments were conducted at the Selçuk University, Veterinary Faculty Dogship Unit. **Gelatinized** starch was found to be 4.87% in pelleted and 17.81% in extruded food ($P < 0.001$). The rate of preference of extruded food by dogs was 0.66. **The pelleted** and extruded dog foods produced in this study were compared in terms of digestibility, fecal score, and cost with the most common one imported and one native product in the market. Dry matter digestibility of pelleted, extruded, imported and domestic commercial dog foods determined by fecal collection method was 81.2%, 84.2%, 83.7%, 83.5% ($P < 0.05$) respectively. **The fecal score** was 3.48 for those who consumed pelleted food, and 3.68 to 3.91 for the others. In calculating the cost, it was determined that extruded food produced in this study could be 5 times more economical than imported food.

Keywords: Dog Food, Pellet, Extrude, Digestibility, Preference

OVEREXPRESSION OF PHAC1 GENE ENCODING PHA SYNTHASE 1 IN PSEUDOMONAS AUREOFACIENS FOR OVERPRODUCTION OF MCL-PHA

EMRE ERDEN^a, SERAP EVRAN^a, NURDAN KASIKARA PAZARLIOGLU^a

^aEge university faculty of science biochemistry department

emrerden@gmail.com

Polyhydroxyalkanoates or PHAs are linear polyesters synthesized and stored in the cell cytoplasm as water-insoluble inclusions by bacteria under the limited nitrogen and excessive carbon sources. These biopolymers are alternative to oil-derived polymers and they have different physical and chemical properties. Besides they are biocompatible and biodegradable, PHAs can be used in most of biomedical applications including tissue engineering as a tissue scaffold, controlled drug release and biomedical implants. **The most common problem** encountered in PHA production is high production costs of PHA. The production costs can be reduced by developing efficient bacterial strains and optimizing the production process. **The aim of** this study is to develop recombinant strain by harboring phaC1 gene encoding PHA synthase 1 to wild-type *P. aureofaciens*. For this purpose, phaC1 gene fragment from *P. aureofaciens* was amplified with PCR and inserted into pBBR1MCS2 plasmid. The recombinant plasmid was firstly introduced into *E. coli* S17-1 and then, it was introduced into *P. aureofaciens* via conjugation in order to carry out the homolog recombination. The efficiencies of recombinant- and wild-type strain were compared. **The results** showed that the mcl-PHA production was increased up to 25% and yield of mcl-PHA production was increased from 4.1% to 7,6% with recombinant strain.

Keywords: PHA Synthase, Overexpression, *P. Aureofaciens*, Mcl-Pha

Acknowledgements: This study was supported by BAP-Ege University/Turkey (2010FEN056).

595

Oral - 14

596 **MODELLING OF PHENAZINE-1-CARBOXYLIC ACID PRODUCTION BY**
597 **PSEUDOMONAS CHLORORAPSIS SUBSP. AUREOFACIENS IN**
598 **SUBMERGED CULTURES AND INVESTIGATION OF ITS**
599 **ANTIMICROBIAL AND ANTICANCER ACTIVITY**

600

KÜBRA ASLAN^a, EMRE ERDEN^a

601

^aEge university faculty of science biochemistry department

602

603

604

cansukubra0735@gmail.com

605 Phenazines and derivatives are nitrogen containing heterocyclic compounds that
606 are synthesized by Pseudomonas and Streptomyces species. These molecules are
607 redox agent with broad spectrum anti-microbial, anti-tumor and anti-parasitic
608 activity. Over the past decades, there has been great interest in their
609 biotechnological production because of their unique properties. **Bacterial**
610 fermentation processes have advantages over chemical synthesis, since necessity of
611 the highly toxic substrates and harsh conditions during the chemical synthesis,
612 formation of toxic by-products and also low productivity. **The aim of the study** is
613 to produce phenazine-1 carboxylic acid (PCA) by bacterial fermentation as a
614 bioactive agent. For this purpose, PCA was produced by P. chlororapsis subsp.
615 aureofaciens on batch mode in submerged cultures. Modelling of PCA production
616 and growth of organism were determined by temporal analysis of variations of
617 biomass, product, depleted carbon and nitrogen source. Chromatographic and
618 spectroscopic methods were used for purification, quantification and
619 characterization of PCA produced by fermentation. Besides, in vitro anti-cancer and
620 antimicrobial activity of characterized pigment was also investigated.

621 **Keywords:** Kinetic Modelling, Phenazine-1-Carboxylic Acid, Anticancer,
622 Antimicrobial, Pseudomonas

623 **Acknowledgements:** *This study was supported by BAP-Ege university/ Turkey (16-Fen-*
624 *005). The authors also would like to thank Ege University ARGEFAR laboratories especially to*
625 *PhD. Güliz AK and MSc Buket Özel for generous support and interpretation in kind of*
626 *cytotoxicity assays.*

627

THE EFFECTS OF SILYMARIN ON THE CELL VIABILITY IN F-98 CELL CULTURE

EFE KURTDEDE^a, REMZI SONER CENGİZ^a, AYŞENUR GÖK^a, UFUK KAYA^b,
GÖRKEM KISMALI^a

^aAnkara university, faculty of veterinary medicine, dept. of biochemistry

^b Ankara university, faculty of veterinary medicine, dept. of biostatic

efekurtdede@gmail.com

Objective: Silymarin is flavonoid has been well studied and reported antioxidant agent. The effects of silymarin on cell viability and proliferations in rat glioblastoma F-98 cell line are investigated. **Methods:** F-98 cells were seeded in 96 well plates in 100 µl medium DMEM. The cells were either left untreated (control) and treated with different concentrations of silymarin (200,100,50,25,12.5,6.2,3.3 µM) for 24, 48, 72 hours. Measurement of silymarin treated and control groups cell proliferation performed with MTT assay and Wound Healing assay. Medium were collected and glucose levels measured to determine cell viability via spectrophotometric method. **Results:** The effects of Silymarin 50-3.3 µM increased cell viability and proliferations in 24, 48 and 72 hours. However significant decrease was observed at 100 and 200 µM in 48 and 72 hours. Difference observed in supernatant levels of glucose activity in 48 hours. (according to control 156,24%, 170%). The increase glucose level of medium indicates that silymarin has cytotoxic effect in F-98 cells at higher concentration. Also in Wound healing assay 100 and 200 µM silymarin according to control 17%, and 0,6% decrease observed in 48 hours. In the MTT assay 100 µM and 200 µM compare with control groups was significant found in 24 and 48 hours. **Conclusion:** These results suggest that role of silymarin dose dependent activity in F-98 cells.

Keywords: Silymarin, F-98, Mtt, Wound Healing

660 **EFFECTS OF INITIAL INOCULUM CULTURE CONCENTRATION ON**
661 **DARK FERMENTATIVE HYDROGEN GAS PRODUCTION FROM**
662 **WASTE PAPER TOWEL HYDROLYSATE**

663 **HIDAYET ARGUN^a, ÖZGÜR GÖKDEMİR^b, İSMAIL GÖRGÜL^c, ZEYNEP SAYLAM^d,**
664 **GÜLİZAR ONARAN^e**

665

666

Pamukkale university

667

668

hargun@pau.edu.tr

669 **This study** reports the effects of initial inoculum culture concentration on dark
670 fermentative hydrogen gas production from waste paper towel hydrolysate. In this
671 context batch dark fermentation experiments were carried out at mesophilic
672 conditions in serum bottles using heat treated anaerobic sludge as inoculum
673 culture. The hydrolysate was obtained after sulphuric acid hydrolysis of waste
674 paper towel and inoculation was carried out after the removal of toxic levels of 5-
675 Hydroxymethylfurfural by resin treatment. The inspected initial inoculum culture
676 concentration interval was between 0-6 g/L and all other factors were kept
677 constant during the experiments. Cumulative hydrogen volumes and hydrogen
678 formation yields were almost the same for all inoculum concentrations. However,
679 specific hydrogen formation rates and lag phases were different. Most effective
680 hydrogen production resulting maximum hydrogen yield (1.11 mol H₂/mol
681 glucose), specific hydrogen formation rate (181.43 mL H₂/ g biomass.h) and total
682 volatile fatty acid formation yield (0.46 g TVFA/g glucose) was obtained at 0.1 g/L
683 inoculum concentration. **This result** was attributed to the most convenient
684 substrate to biomass ratio (0.014 g biomass/g substrate) at 0.1 g/L initial
685 inoculum concentration.

686

687 **Keywords:** Waste Paper Towel, Hydrogen Production, Acid Hydrolysis, Initial
688 Inoculum Concentration

689

HYDROGEN GAS PRODUCTION USING HOT SPRING MICROFLORA

HIDAYET ARGUN^a, AYŞE ÇALKAYA^b, HANDAN HAN^c, İKBAL ÖREN^d

ZEYNEP KARAOĞLAN^e

Pamukkale university

hargun@pau.edu.tr

Hydrogen (H₂) is considered as one of the most attractive fuels of the future.

However, most of the H₂ in the market is produced from natural gas. In this

context H₂ production from waste materials has gained attention due to the

depletion risk of fossil fuel reserves in the future and environmental concerns. In

this study H₂ was produced from molasses using hot-spring culture as inoculum.

Highest H₂ yields in the literature are reported at thermophilic conditions and

therefore the hot-spring was supposed to be a source for thermophilic H₂

producing microorganisms. Hence H₂ production at psychrophilic, mesophilic and

thermophilic conditions were tested at different inoculations. Among three different

tested temperature scales mesophilic conditions with ten percent inoculum

addition (v/v) resulted most efficient H₂ formation. **The hot-spring culture** was

found to be an effective inoculum source for H₂ production from molasses.

Keywords: Molasses, Dark Fermentation, Hydrogen, Hot-Spring

714 **A NEW HUMANIZED MOUSE MODEL FOR AUTO-IMMUNE**
715 **MYOCARDITIS AND ITS USE TO DEVISE IMMUNOMODULATION**
716 **THERAPY**

717 **MEHMET EMRAH ŞELLI^a, DAVID C WRAITH^b, ANDREW C NEWBY^c**

718

^aUşak university

719

^bUniversity of birmingham

720

^cUniversity of bristol

721

722

723

emrah.sell@usak.edu.tr

724 **Myocarditis** is the principal cause of heart failure in young adults. Its progression
725 to dilated cardiomyopathy is associated with the development of auto-immunity,
726 especially to human cardiac α -myosin (hCAM). Consistent with this, HLA genotype
727 influences prevalence of the disease. Previous studies showed that humanised DQ8
728 transgenic non-obese diabetic mice spontaneously developed autoimmune
729 cardiomyopathy, whereas the DR4 allele is over represented in patients and there is
730 no association with diabetes. We therefore attempted to induce experimental
731 autoimmune myocarditis in DR4 transgenic mice (DR4 mice) as a more relevant
732 model of the human disease. **DR4 mice** were injected with purified hCAM or vehicle
733 subcutaneously in complete Freund's adjuvant (CFA). After 3 weeks, anaesthetised
734 mice were subjected to cardiac ultrasonography. The hearts were then perfused
735 fixed for histology and spleens were harvested for proliferation assay. *in silico*
736 predicted potentially immunomodulatory peptides were proven water soluble and
737 effective in T-cell proliferation assays. For immunotherapy, mice were pre-dosed
738 with escalating doses of mixtures of 3 each of 6 soluble hCAM-derived peptides
739 (pools 1 and 2) according to an established protocol. **All mice** immunized with
740 hCAM developed high titres of both IgG1 and IgG2c antibodies. Consistent with
741 this, splenic T-cell proliferation responses to hCAM significantly increased
742 compared to un-immunized mice. Immunized mice showed a significant decline in
743 cardiac output and fractional shortening and increase in diastolic dimension. 5/5
744 immunized vs 0/5 control mice showed cardiac inflammation based on histology.
745 Pre-treatment with hCAM derived peptide pools 1 or 2 blunted the T-cell
746 proliferation response and pool 2 also decreased both IgG1 and IgG2c levels. Pools
747 1 and 2 significantly improved the left ventricular cardiac. Pool 2 also significantly
748 reduced cardiac inflammation. **We have developed a novel**, more relevant
749 humanized mouse model of autoimmune myocarditis and demonstrated its ability
750 to validate the immunomodulatory activity of hCAM derived peptides.

751 **Keywords:** Autoimmunity, Myocarditis, Cardiomyopathy, Immunotherapy

AUTOIMMUNITY TO HSP60 DURING DIET INDUCED OBESITY IN MICE

MEHMET EMRAH ŞELLI^a, DAVID C WRAITH^b, ANDREW C NEWBY^c

^aUşak university

^bUniversity of birmingham

^cUniversity of bristol

emrah.sell@usak.edu.tr

Although the association of a chronic low-grade inflammation with obesity has long been appreciated, its molecular basis is yet to be defined. The proven involvement of adaptive immunity, coupled with a phenotypic switch from autoimmune suppressive tolerogenic Treg to pro-inflammatory CD4⁺ Th1 and CD8⁺ T cells, during progression of obesity necessitates the presence of a triggering antigen as an activator of T and B cells. **HSP60** is an evolutionary conserved mitochondrial chaperonin which can be located in the cytosol or exposed on the cell membrane also. An increase in cell membrane HSP60, which may be accompanied by HSP60 release into circulation, is especially considered a signal of autoimmunity. HSP60 is linked to obesity as a mediator of adipose tissue inflammation and insulin resistance. Moreover, circulating HSP60 levels are found to be higher in obese individuals than lean controls (Märker et al. 2012). **We observed** an increase in circulating HSP60 levels and an adaptive immune response against HSP60 at both T cell and B cell (antibody) levels during continuous high fat feeding of C57bl6 mice. Hence HSP60 appears to be one of the mystery auto-antigens triggering the early T and B cell responses during obesity. Furthermore, we attempted a peptide therapy in a dose escalation protocol aiming to down-regulate the inflammatory related adverse effects of obesity by achieving tolerance in T cell populations and suppressing the pathogenic antibody response. Treatment with a mixture of three proven immunomodulatory HSP60 peptides did not reduce weight but completely reversed the increase in VLDL/LDL levels and partially reversed the glucose intolerance in obese mice, which encourages further research to improve peptide therapy. **We clearly showed** that obese mice mount an autoimmune response to HSP60, which partly underlies the resulting metabolic disturbances.

Keywords: Autoimmunity, Auto-Antigen, Obesity, Immunotherapy

INVESTIGATION OF BIOSORPTION KINETICS OF AN AGRICULTURAL RESIDUAL ON DYE REMOVAL

LEVENT GÜREL

Pamukkale university

lgurel@pau.edu.tr

This study was conducted to examine biosorption kinetics of an agricultural residue called pepper seeds. Several adsorption models were applied to the results obtained from the kinetic tests of pepper seed biosorbent on dye solutions. In many literature papers, the most used models for kinetic analyses were pseudo-first and pseudo-second order kinetic models. So to evaluate the other models on kinetic analyses, detailed studies were conducted using different kinetic models. The kinetic adsorption models used in this work are Elovich, Weber-Morris, liquid film diffusion and Dumwald-Wagner kinetic models. At the end of these kinetic analyses, it was found that the Dumwald-Wagner kinetic model best described the biosorption of dye by using this biomass. The R-squared (R²) results obtained for the kinetic models were found as 97.75, 95.48, 94.18 and 99.30% for the kinetic models of Elovich, Weber-Morris, liquid film diffusion and Dumwald-Wagner, respectively. High R-squared (R²) values obtained in this study for different kinetic models along the mostly used pseudo-first and pseudo-second order kinetics showed that all these kinetic models should be calculated and evaluated in all studies regarding biosorption processes to ensure the biosorption mechanism.

Keywords: Biosorption Kinetics, Dye Removal, Pepper Seed, Wastewater Treatment

817

Oral - 21

818

INVESTIGATION OF NICKEL BIOSORPTION BY AN AGRICULTURAL RESIDUAL

819

820

LEVENT GÜREL^a, TUĞBA MEŞE^b, ARIFE ŞEKER^c, ASLI KILIÇ^d, ALI SERKAN

821

YILDIRIM^e

822

823

^aPamukkale university

824

825

lgurel@pau.edu.tr

826

The removal of an important heavy metal found in water bodies because of
improper discharge from several industries was investigated in this study by using
an agricultural residual material called pepper seed. Peppers are very important
foodstuff for Turkey. Especially, they are extremely used in several industrial plants
to obtain pepper paste. Also, paste manufacturing in houses is very widespread in
some regions of the country. The seeds of the peppers are the residuals, and they
are generally thrown away from houses and also industries. So, to evaluate the
treatment potential of these residuals, biosorption studies were conducted. Raw
pepper seeds with no chemical pre-treatment were used. The pepper seeds used in
these studies were in the size of 125 µm. Optimum pH for the tests were found as
5.5 in pH studies. Also, a maximum sorption capacity of 6.33 mg/g was obtained in
all studies carried out for nickel removal. **At the end of the results**, it was seen
that this residual material had a potential for nickel removal from solutions.

839

840

Keywords: Nickel Removal, Wastewater Treatment, Biosorption, Pepper Seed

841

843 **ASSOCIATION OF MNSOD GENE POLYMORPHISM AND SOD**
844 **ACTIVITY AND THE RISK OF CORONARY ARTERY DISEASE IN**
845 **PATIENTS WITH NON-DIABETIC METABOLIC SYNDROME**

846 **SEDA GÜLEÇ YILMAZ^a, ATIKE TEKELI KUNT^b, ORHAN FINDIK^c, SELİM İSBİR^d,**
847 **TURGAY İSBİR^e**

848

849 ^aDept. of molecular medicine, institute of health sciences, Yeditepe uni., İstanbul, Turkey

850 ^bDept. of cardiovascular surgery, Ankara numune education and research hospital, Turkey

851 ^cDept. of cardiovascular surgery, Derince education and research hospital, Kocaeli, Turkey

852 ^dDept. of cardiovascular surgery, Marmara university, faculty of medicine, İstanbul, Turkey

853 ^eDept. of medical biology, faculty of medicine, Yeditepe university, İstanbul, Turkey

854

855 seda0802@yahoo.com

856 **BACKGROUND AND AIM:** The purpose of this study was to investigate the activity
857 of MnSOD and also analyze the association of MnSOD gene polymorphism and the
858 risk of CAD in a cross-sectional study of non-diabetic MetS patients. **METHODS:** A
859 total of 81 adult subjects manifesting metabolic syndrome (MetS) participated in
860 this study. The non-diabetic MetS patients were divided into two groups either
861 having coronary artery disease (CAD) (n= 29) or not (n= 36). Blood samples were
862 stored in EDTA tubes and DNA samples were extracted from whole blood with a
863 salting-out procedure. **RESULTS:** Statistical analysis revealed no association of
864 genotype on the risk of CAD development (OR= 1.180, 95% CI= 0.542-2.570, p=
865 0.676). However MnSOD genotype was not associated with plasma levels of
866 MnSOD activity (p= 0.468, mitochondrial MnSOD activity was shown to be lower
867 in CAD (+) than the CAD (-) group (p= 0.036). Also MnSOD activity was reported to
868 be lower in male population than the females (p=0.003) and CAD risk was more
869 common in males than females (p=0.007). **CONCLUSIONS:** We have demonstrated
870 that lowest SOD activity was associated with increased risk of CAD and male
871 gender was associated with lowest SOD activity and also increased risk of CAD in
872 this non-diabetic MetS population. However we could not demonstrate an
873 association with CAD and MnSOD activity with MnSOD Ala16Val polymorphism.
874 Further prospective randomized studies with larger number of patients are
875 warranted to confirm or refute these results.

876

877 **Keywords:** Mnsod, Polymorphism, Metabolic Syndrome

879 **COMPARISON OF THE BENEFICIAL EFFECTS OF VITAMIN E AND N-**
880 **ACETHYL CYSTEINE SUPPLEMENTATIONS WITH THERAPEUTIC**
881 **ULTRASOUND APPLICATION ON THE TREATMENT OF MUSCLE**
882 **INJURY OF RATS**

883 **AYSE MERVE^a, HALIL OZKOL^b, YASIN TULUCE^c**

884

885 ^aDept. of physiotherapy and rehabilitation, faculty of health science, Necmettin Erbakan
886 university

887 ^bDept. of medical biology, faculty of medicine, Yuzuncu Yıl university

888 ^cDept. of medical biology, faculty of medicine, Yüzüncü Yıl university

889

890

amervetat@gmail.com

891 **Background:** Increased oxidative stress has been reported after muscle injury.

892 **Objective:** In this study, for the first time in the literature, ameliorative efficiency of
893 some antioxidant substances such as vitamin E, N-acetyl cysteine were compared
894 with therapeutic ultrasound application in muscular injury of rats. **Method:** Thirty-
895 six rats were equally divided into 6 groups: control (C), only injury (OI), therapeutic
896 ultrasound (U), vitamin E (Vit E), N-acetyl cysteine (N) and mixed (M). Serum
897 creatine kinase (CK), aspartate aminotransferase (AST), alanine aminotransferase
898 (ALT), lactate dehydrogenase (LDH), interleukin-6, TNF- α levels and whole blood
899 myeloperoxidase (MPO) activity were measured in laboratory. **Results:** Significant
900 ($p < 0.05$) elevations of CK, AST and ALT as well as slight ($p > 0.05$) escalation of LDH
901 were observed in OI group compared with C group. Activities of CK, LDH and AST
902 reversed to C values in almost all treatment groups ($p > 0.05$), whereas ALT
903 decreased markedly ($p < 0.05$) only in Vit E group. When comparing with C, MPO
904 activity increased in OI and significantly ($p < 0.05$) decreased in all treatment groups
905 (except U). With regard to serum interleukin-6 and TNF- α , slight ($p > 0.05$) increases
906 of their levels were observed in all groups compared to C. **Conclusions:** Our data
907 revealed that supplementations of Vit E and NAC with TU application may be
908 beneficial in treatment of muscle injury. Patient with muscle injury should be
909 encouraged in vitamin and antioxidant intake.

910

911 **Keywords:** Muscle Injury, Oxidative Stress, Therapeutic Ultrasound, Antioxidant.

912

913
914 **INVESTIGATION OF BIOCHEMICAL AND HISTOPATHOLOGICAL**
915 **EFFECTS OF VITAMIN C, SELENIUM AND THERAPEUTICAL**
916 **ULTRASOUND APPLICATION ON MUSCULAR INJURY FORMED IN**
917 **RATS**

918 **NECATI MUHAMMED TAT^a, YASIN TULUCE^b, MEHMET KARA^c**
919 **AYSE MERVE TAT^a, HALIL OZKOL^b**

920
921 ^a Necmettin Erbakan university, health sciences faculty, dept. of physiotherapy and
922 rehabilitation

923 ^b Yüzüncü Yıl university faculty of medicine department of medical biology

924 ^c Yüzüncü Yıl university faculty of medicine department of physiology

925
926 necatimuhammedtat@gmail.com
927

928 **INTRODUCTION:** Muscle injury is the type of soft tissue injury and then occurred
929 increased oxidative damage . **PURPOSE:** The efficiency of therapeutic ultrasound
930 treatment and various antioxidant substances was compared in experimental
931 muscular injury. Some serum enzymes, oxidative stress biomarkers,
932 myeloperoxidase (MPO) activity, inflammation markers interleukin-6 (IL-6) and
933 tumor necrosis factor- α (TNF- α) were evaluated histopathological examination.
934 **METHODS:** Each group enclosing 6 rats were constituted to six groups control,
935 only injury (OI), ultrasound (U), vitamin (Vit C), selenium (S) and mix (M). The
936 muscle injury was formed through laceration of gastrocnemius muscle. Treatment
937 was applied three times to groups. Therapeutic ultrasound, vitamin C, selenium
938 and combination of them was applied respectively to the rats. Rats were sacrificed
939 in the sixth day. Blood and tissue samples were obtained for biochemical and
940 histological analyses. **RESULTS:** While OI group compared to control in terms of
941 serum enzymes, significant ($p < 0.05$) elevations of CK, AST and ALT as well as slight
942 increase of LDH. CK, ALT and AST levels reversed to control values in almost all
943 treatment groups ($p < 0.05$) as well as slight decrease of LDH. The shifts in total
944 antioxidant level were insignificant. Total oxidant status (TOS) and oxidative stress
945 index (OSI) increased in OI group and it decreased in S and M. However; among
946 these alterations only TOS in muscle tissue was found to be statistically significant
947 ($p < 0.05$). While MPO activity increased in OI group, it decreased other treatment
948 groups. However only the increase on MPO in the blood tissues of OI group and the
949 decrease in Vit C group were statistically significant ($p < 0.05$). There were not any
950 marked alteration between groups in terms of serum inflammation markers (IL-6
951 and TNF- α) and histological findings. **CONCLUSION:** Results of this study revealed
952 that apart from therapeutic ultrasound treatment, supplementation of vitamin C
953 and selenium may also be useful treatment of muscular injury. However further
954 studies are needed to support these results.

955 **Keywords:** Rats, Muscular Injury, Therapeutic Ultrasound, Vitamin C, Selenium,
956 Oxidative Stress.

THE EFFECT OF ANKAFERD BLOOD STOPPER AND POLYVINYL PYRROLIDONE ON THE MODEL ORGANISM

MEHMET OKAN ERDAL^a, EDA GÜNEŞ^b, LOKMAN GEMİ^c

^aNecmettin Erbakan niversity meram vocational school

^bNecmettin Erbakan university, faculty of tourism, dept. of gastronomy and culinary arts,

^cNecmettin Erbakan university meram vocational school department of food processing

moerdal@konya.edu.tr

Polyvinyl pyrrolidone (PVP) which is soluble in organic liquids similar to proteins, is used in pharmaceuticals and many fields (food, beauty, cleaning products and medical applications, photography, testing, glue making). Ankaferd Blood Stopper (ABS) is used as a haemostatic agent. Despite their useful potential, nanofiber PVP and ABS containing nanofibers have not been studied for their potential for co-use, and in this study their availability for ecosystem and health has been investigated. For this purpose diets were coated with 10% PVP (produced by the electrospinning, ABS (0,1-1%) and PVP-ABS. *Drosophila melanogaster* Meigen larvae were feed and matured with these diets survival and sex ratio, development time were determined. It has been determined that the separate use of the two substances (PVP and ABS 1%) were reduced the proportion of mature and female individuals. The combined use of PVP and ABS were found to have a positive effect, similar to the control. **As a result**, the nanofiber and drug (ABS 1%) can be used in the medical field, but more detailed studies are needed to make definite inferences

Keywords: Nanofiber, Polyvinyl Pyrrolidone, Ankaferd, *Drosophila Melanogaster*, Nutrition

INVESTIGATION OF PARVOVIRUS B19 IGM AND IGG POSITIVITY RATES IN PEDIATRIC HEMATOLOGY PATIENTS

AYSUN GÖRKEM^a, AYŞE RÜVEYDA UĞUR^b, FATMA TAŞBENT^c, MEHMET

ÖZDEMİR^d, MAHMUT BAYKAN^e

^aDept. of medical microbiology, faculty of meram medicine, Necmettin Erbakan university,
Konya, TURKEY

mbaykantr@hotmail.com

Abstract:

996 Human parvovirus B19 is a frequent etiologic agent causing erythema infectiosum
997 in children. It has recently been suggested that parvovirus B19 may be latent after
998 infection and cause reactive infections especially in immunosuppressive patients
999 with hematological problems. In this study, we aimed to investigate the parvovirus
1000 B19 IgM and IgG positivity rates in sera of patients consulted Pediatric Hematology
1001 clinic. **We retrospectively** screened the laboratory results of parvovirus B19 IgM
1002 and IgG antibody assays of children < 18 years, who consulted Pediatric in and
1003 outpatient clinics at between 2013-2016. Parvovirus B19 IgM and IgG antibodies
1004 were investigated in serum samples by using enzyme linked immunosorbent assay
1005 method in the Medical Microbiology Laboratory. **Regarding** all kinds of pediatric in
1006 and outpatients, the total number of patients from whom parvovirus B19 IgM
1007 antibodies were investigated was 2697 and parvovirus B19 IgM antibodies were
1008 detected in 168 (6.2%). On the other hand, of all, the number of patients attending
1009 Pediatric Hematology clinics was 602 and parvovirus B19 IgM antibodies were
1010 detected in 109 (18.1%). A total of 1299 patients were asked for parvovirus B19 IgG
1011 antibody and 368 (28.3%) of them were positive for parvovirus B19 IgG. Of all,
1012 parvovirus B19 IgG antibody was detected in 244 (25.6%) of the 952 samples
1013 collected from Pediatric Hematology clinics. Parvovirus B19 IgM and IgG positivity
1014 in samples from Pediatric in and outpatient clinics other than Pediatric Hematology
1015 were 2.8% and 35.7%, respectively. Parvovirus IgM positivity in serum samples
1016 sent from the Pediatric Hematology in and outpatients was statistically significant
1017 compared to those sent from Pediatric clinics other than Pediatric Hematology (p =
1018 0.0001). **The higher detection** rate of serum parvovirus B19 IgM positivity in
1019 patients under the follow-up of Pediatric Hematology clinics suggests that immune
1020 suppression-related virus reactivation or persistence may occur in these patients.

Keywords: Parvovirus B19, Igm, Igg, Child Patient

1023

Oral - 27

1024

EFFECT OF BLOOD HORMONE LEVELS AND BODY CONDITION SCORE IN SUCCESS OF ARTIFICIAL INSEMINATION IN COWS

1025

1026

İLKER YAVAŞ^a, İSHAK GÖKÇEK^b, ECE KOLDAŞ^c, TUĞBA KORKMAZ YAVAŞ^d

1027

1028

^aMustafa Kemal üniversitesi veteriner fakültesi dölerme ve suni tohumlama a.d

1029

^bMustafa Kemal üniversitesi veteriner fakültesi fizyoloji a.d

1030

^cMustafa Kemal üniversitesi veteriner fakültesi doğum ve jinekoloji a.d

1031

^dHatay il tarım müdürlüğü gıda kontrol laboratuari

1032

1033

ishakgokcek@hotmail.com

1034

Abstract:

1035

Fertility rate with artificial insemination obtained depends on various factors. One of these factors is the body condition score. In this study, the body condition scores determined during artificial insemination, follicle development, steroid hormone (estrogen) and the impact on the level of estrus symptoms and aimed to reveal the possible cause of infertility. **In this study**, cows with no obstacles impeding fertility and at least one birth without any problems have been used at the private farms in Hatay Region. **Body Condition Score** : The oestrus assessments of cows that have been identified have been made according to the following criteria: Uterian flow, edema of Vulva, Vagina's color, Duldung reflex and diameter of follicle. **Cows** that are determined to be at estrus were inseminated by recto-vaginal method. **Estrogen** levels were determined by immunological methods (ELISA). **As a result**, it has been observed that if the body condition score is less than 2.5 in artificial insemination, the amount of estrogen is decreased, the first insemination pregnancy rate may decrease from 10% to 18% and the birth-first insemination interval may be prolonged.

1050

1051

Keywords: Body Condition Score, Infertility, Hormone Levels, Artificial Insemination.

1052

1053

Acknowledgements: *This research was financed by Research Fund of Mustafa Kemal University (Project Number:13882)*

1054

1055

1057 **EFFECTS OF ENTODINIUM CAUDATUM MONOCULTURE**
1058 **INOCULATION ON IN VITRO FERMENTATION, METHANE**
1059 **PRODUCTION AND PREVENTION OF SUB-ACUTE RUMINAL**
1060 **ACIDOSIS**

1061 **NURETTIN GÜLŞEN^a, HUZUR DERYA ARIK^a, MUSTAFA SELÇUK ALATAŞ^a,**
1062 **MUHAMMAD NAEEM TAHIR^b**

1063

1064 ^aDept. of animal nutrition and nutritional disorders faculty of veterinary medicine, Selcuk
1065 university, Konya, TÜRKİYE

1066 ^bUniversity college of veterinary and animal sciences, the islamia university of bahawalpur

1067

1068

nurettingulsen@gmail.com

1069 The **objective** with this study was to look at the possibilities to reduce the chances
1070 of sub-acute ruminal acidosis (SARA) by the use of protozoa (Entodinium
1071 caudatum) monoculture. This study included the in vitro fermentation and gas
1072 production recordings of two substrates i.e. wheat and corn incubated with two
1073 varying levels (1 and 2 ml of monoculture with 5.5x10⁴/ml) of E. caudatum. The
1074 rumen fluid used to study was supplied by 2 rumen cannulated holstein heifers
1075 (450 kg live weight) fed with alfalfa only. 30 g/L wheat and 50 g/L maize substrate
1076 were used to induce SARA in vitro. The fermentation bottles contained the
1077 substrate, incubation medium, protozoan free rumen fluid and E. caudatum
1078 monocultures and incubations continued for 24 hours. From the beginning of the
1079 incubation, it was observed that the protozoa were motile and started to engulf the
1080 substrate. The mobility was reduced after eight h of incubations as the inside was
1081 completely filled with substrate. The concentrations of volatile fatty acids (VFA) and
1082 lactate as well as general fermentation parameters did not change in the medium
1083 as a result of E. caudatum inoculation or substrate. However, E. caudatum
1084 inoculation increased the cumulative gas production (P <0.002) and tended to
1085 increase the methane concentration in the total gas (P <0.061). **It is concluded**
1086 that E. caudatum monoculture inoculation have little importance in reducing the
1087 chances of SARA as a result of poor relationship between protozoa and lactate
1088 levels in the rumen in vitro. **Additionally**, it may enhance methane production in
1089 the total gases produced in the rumen, thus posing serious impacts on climate

1090 **Keywords:** Sub-Acute Ruminal Acidosis, In Vitro Rumen, Entodinium Caudatum,
1091 Methane

EFFECTS OF METOCLOPRAMIDE IN MATERNAL RAT BRAIN: USE TO INCREASE OF BREAST MILK PRODUCTION

OYA SAYIN^a, SEREN GÜLŞEN GÜRGEN^b

^aDokuz eylul university faculty of medicine, research laboratory, İzmir, TURKEY,

^bDept. of histology and embryology, school of vocational health service, Celal Bayar university, Manisa, TURKEY.

oya.sayin@deu.edu.tr

Background: Metoclopramide is an anti-emetic drug widely used in nausea and vomiting, and also has recently used for increasing maternal breast milk production in the lactation period. **Purpose:** The aim of this study was to investigate the extrapyramidal effects of metoclopramide to induce breast milk output in the lactation period in the central nervous system of mother. **Methods:** 18 young adult female Wistar albino rats that had recently given birth, together with their pups, were divided into 3 groups; Group 1: healthy control. Group 2: administered a low dose of MCP (10 mg/kg for 21 days, twice daily, ip), and Group 3: administered a high dose of MCP (45 mg/kg 21 days, twice daily, ip). The experiment continued throughout the lactation period. Each mother's brain tissues and blood samples were removed at the end of the 21st day. Dopamine D2 receptor (DRD2), brain derived neurotrophic factor (BDNF) and neural growth factor (NGF), markers of extrapyramidal reaction in the brain, as signal molecules were assessed using the immunohistochemical staining. DRD2, BDNF and NGF levels were performed using the ELISA kits in the brain tissues and blood samples. **Results:** Immunohistochemical and biochemical evaluations showed that there were no statistical difference in expression and serum levels of DRD2, BDNF and NGF between the three groups ($p < 0.05$). **Conclusions:** The results of this study suggest that there were no differences between three groups regarding expression and serum levels of DRD2, BDNF and NGF. The metoclopramide no have effect on the mother, on the other hand, our previous study, we show that metoclopramide causes the extrapyramidal reactions in newborn brain tissue. For this purpose, metoclopramide should be used carefully during in the lactation period by mothers.

Keywords: Metoclopramide, Brain, Extrapyramidal Reactions

1127

Oral - 30

1128

THE STUDY OF THE EFFECT ON THE BONE TISSUE DEPENDING ON THE AGE GINGKO BILOBA EXTRACT AND GROWTH FACTORS

1129

1130

SEREN GÜLŞEN GÜRGEN^a, GÜLCE NAZ YAZICI^b

1131

1132

^amanisa celal bayar university

1133

^bberzincan university

1134

1135

serengurgen@gmail.com

1136

AIM: Ginkgo Biloba leaf extract is a medicine used to treat Alzheimer, vascular demance and mental memory disorders. Aim of this study the effect of Ginkgo biloba extract on bone tissue and the epiphyseal cartilage is investigated in comparison between groups of young and old in terms of apoptotic factors.

1137

1138

1139

1140

MATERIALS-METHODS: In the study, 40 male wistar albino rats were divided into 4 groups. Group 1. 30 days young control (serum physiological, 2 months, 2 doses) Group 2: 30 days young ginkgo (GbE 100 mg/kg/days, 2 months, 2 doses) Group 3. 24 months old control (serum physiological, 2 months, 2 doses) Group 4:

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

RESULTS: FGF-2 expression in the epiphyseal plate of young ginkgo and young control groups, particularly around the ossification in the cartilage matrix in proliferating chondrocytes was quite strong. In addition, the young control group was observed in osteoblasts and endosteum FGF-2 immunoreaction is moderate young ginkgo groups was strong release. FGF-2 expression was observed stronger in the elderly proliferation in the control group, the old ginkgo group was remarkable reduction in the reaction. In both groups, FGF-2 release is weak identified as ossification areas. In addition, osteoblasts, FGF-2 reaction is quite different from the young group decreased. PDGF-A immunostaining in all groups were weak to moderate staining in tissue.

1157

1158

1159

1160

CONCLUSIONS: Ginkgo biloba aged younger than those in the bone tissue, especially in chondrocytes FGF-2 expression in osteoblast cells, and further increase in proliferation. For this reason, bone structure than those of the elderly than the young group concluded that a positive effect. Development of bone formation and release of PDGF-A to have no effect on the FGF-2.

1161

1162

Keywords: Ginkgo Biloba, Bone, FGF-2, PDGF-A

1163

1164

Oral - 31

1165

D-LACTIC ACID PRODUCTION FROM CORN COB USING L-LDH GENE DEFICIENT LACTOBACILLUS HELVETICUS OZH12

1166

1167

HARUN ÖNLÜ^a, SAADET BALTACI^b, HALUK HAMAMCI^c,

1168

OZLEM OSMANAGAOGLU^d

1169

1170

^aMuş Alparslan university

1171

^bMiddle east technical university, department of food engineering, 06800 Ankara, TURKEY

1172

^cMiddle east technical university, department of food engineering, 06800 Ankara, TURKEY

1173

^dAnkara university, faculty of sciences, department of biology, biotechnology units

1174

1175

harunonlu@gmail.com

1176

Introduction: In recent years, concerns over the microbial production of commercially valuable products have been growing. Lactic acid has great potential for the production of biodegradable and biocompatible poly-lactic acid (PLA) polymers that drive the current market expansion for lactic acid which is a good alternative to petroleum-based plastics. Lactic acid is a chiral molecule that present in nature as two different optical isomers (D-LA and L-LA). D-Lactic acid is used for the production of poly d-lactic acid (PDLA). These pure polymers (PLLA and PDLA) are relatively heat sensitive, while stereocomplexes of PLLA and PDLA have a melting point ~50 °C higher than their respective pure polymers and are more biodegradable. The ratio of L- and D-lactic acids influences the properties and the degradability of PLA. These findings increased the interest to D lactic acid.

1187

Method: In this study, in order to meet the demand of the existing D-lactic acid, L-lactate dehydrogenase gene of Lactobacillus helveticus OZH12 was knock-out and disruption and substitution of the ldh-L gene of OZH12 were carried out using pGIT707 plasmid-based double-crossover homologous integration. We designed a fermentation process using corn cob which is industrial waste for production of lactic acid. Total lactic acid and L-lactic acid production were measured by HPLC and Lactate analyzer, respectively. D-lactic acid production was calculated by subtracting L-lactic acid difference from total lactic acid. **Results and Discussion:** The results showed that clone P8 consumed 41% of the available sugar, and produced totally 25.5 mg / ml lactic acid. 22,6 % of this lactic acid is D-lactic acid and 2,9 % is L-lactic acid. These findings indicate that the strain can be used in the production of lactic acid by using industrial wastes.

1199

1200

Keywords: L-Ldh Gene, Lactic Acid Bacteria, Lactic Acid Production, Knock-Out, Corn Cob

1201

1202

Oral - 32

1203

PROTECTIVE EFFECT OF OMEGA-3 FATTY ACIDS ON MSG INDUCED HIPPOCAMPAL NEURONAL APOPTOSIS IN THE ADOLESCENT RATS

1204

1205

NURCAN UMUR^a, S. GÜLŞEN GÜRGEN^b

1206

^aManisa Celal Bayar university, vocational school of health services, molecular biology

^bManisa Celal Bayar university, vocational school of health services, histology-embryology

1209

1210

nurcanumur@gmail.com

1211

Monosodium glutamate (MSG) is used as a flavor enhancer in processed foods and was reported to have neurotoxic effects. The aim of this study is to investigate the possible toxic effect of MSG on histological and apoptotic features of hippocampus of adolescent rats to evaluate the possible protective role of eicosapentaenoic (EPA) and docosahexaenoic acid (DHA) against this effect. **Wistar** albino rats divided 5 groups 6 adolescent in each one. 1 Group: Healthy control. 2. Group: MSG applied group (4 mg/kg 1.3.5.7.9. days intraperitoneal (IP)), 3. Group: MSG+ EPA applied group (MSG + 300 mg/kg 9 days IP), 4. Group: MSG + DHA applied group (MSG + 300 mg/kg 9 days IP), 5. Group: MSG + EPA + DHA applied group (MSG + 300+300 mg/kg 9 days IP), 9 days later brain tissues took from rats. TUNEL technique was performed. Examination of the hippocampus of adolescent brain revealed powerful TUNEL positive reaction in granular form in the neuron cytoplasm of the CA1 and DG areas in the MSG group. Expression weak in both areas in the control group. Decreased reaction was determined in hippocampal neurons in all areas in the MSG-EPA, MSG-DHA and MSG-EPA+DHA groups. No significant difference was observed in the expression between the MSG-EPA and DHA groups. We **conclude** that since MSG caused an increase in TUNEL positive expression in hippocampus CA1 an DG area in the brain, while MSG-EPA and MSG-DHA caused a decreased apoptosis in hippocampal neurons from both CA1 and DG areas, care needs to be taken with the omega-3 fatty acids use of MSG in the adolescent period due to the possibility of neuronal apoptosis.

1232

Keywords: Msg, Epa, Dha, Brain, Tunel

1234

A DEEP CONVOLUTIONAL NEURAL NETWORK MODEL FOR DETECTING DIABETIC RETINOPATHY

BUSE MELIS ÖZYILDIRIM^a, MUSTAFA ORAL^b, SERKAN KARTAL^c

^aComputer engineering,engineering and architecture faculty,Çukurova university

^bComputer engineering,engineering and architecture faculty,Çukurova university

^cComputer engineering,engineering and architecture faculty,Çukurova university

e

mozyildirim@cu.edu.tr

In this study, a deep convolutional model is proposed and its performance is compared with that of transfer learning methods such as GoogleNet, AlexNet. There exist some deep learning studies in the literature for detecting diabetic retinopathy. Generally, Kaggle dataset which classifies images into five categories according to the level of disease, was utilized in these studies. Unlike other studies existing in the literature, in this study lesions (hemorrhagic, soft and hard exudates, red dots) are detected separately. DIARETDB1 and DIARETDB0 datasets were utilized for training and testing the proposed model. Firstly, these images were cropped into 129x129 size images and these images were labelled according to the lesion types that they include. Then, for each lesion, different convolutional model was trained. Finally, according to the results from each trained model, a report was provided. 5-cross validation was implemented to test the proposed model, GoogleNet, and AlexNet. **According to the test results**, while proposed model achieved almost 80% overall classification accuracy, GoogleNet and AlexNet provided almost 60% and 47%, respectively.

Keywords: Diabetic Retinopathy, Deep Learning, Convolutional Neural Network

1263

1264

SHEAR BOND STRENGTH OF CEROMER BONDED TO ZIRCONIUM OXIDE INFRASTRUCTURE TREATED WITH DIFFERENT TYPES OF LASERS

1265

1266

1267

**OGUZHAN GORLER^a, IHSAN HUBBEZOGLU^b, MELIH ULGEY^c, RECAI ZAN^d,
SERRA KUTLU^e**

1268

1269

^aCumhuriyet university faculty of dentistry

1270

1271

1272

drgrlr@ymail.com

1273

Objectives. Laser modalities are investigated for the preparedness of infrastructures for the bonding of superstructures. We aimed to examine the shear bond strength (SBS) of ceromer and nanohybrid composite to zirconium oxide (ZrO₂) infrastructures treated with Er:YAG, Nd:YAG, and KTP laser modalities in vitro settings. **Research Methods.** The study specimens had 2 sets including 32 ZrO₂ infrastructures with ceromer superstructures and 32 ZrO₂ infrastructures with nanohybrid composite superstructures. These specimens divided randomly into 4 treatment modalities (n=8): no treatment (controls), Er:YAG, Nd:YAG, and KTP lasers. The ZrO₂ infrastructures were prepared in the final dimensions of 7 mm in diameter and 3 mm in thickness in line with ISO 11405 standard. Ceromer as superstructure was applied after their surface treatments according to selected treatment modality. SBS test was performed to test the preparedness of ZrO₂ infrastructure to bonding obtained after laser preparation. **Results.** All laser preparations had positive effect on the SBS value compared to no treatment, and with ceromer and ZrO₂ combination, Nd:YAG and KTP laser preparations were more effective than Er:YAG laser preparation; and with nanohybrid composite and ZrO₂ combination, Nd:YAG, Er:YAG, and KTP laser, in order of decreasing efficacy, had positive effect on the SBS value. When the SBS values of ceromer and nanohybrid composite were compared, there were significant differences among specimens prepared with Er:YAG, Nd:YAG, and KTP lasers and no treatment (p<0.05). In all preparation modalities, the nanohybrid composite specimens had more SBS value compared to the ceromer ones (p<0.05) except KTP laser preparation that was more effective in ceromer specimens (p<0.05). **Conclusions.** Overall, in current experimental settings, KTP, Nd:YAG and Er:YAG lasers, in order of strength, increases the bonding of ceromer to ZrO₂ infrastructure. The studied laser modalities need to be subjected for further studies to determine their appropriate settings and specifications for their future usage in prosthodontics.

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

Keywords: Er:YAG Laser, Nd:YAG Laser, KTP Laser, Ceromer, Zirconium Oxide, Shear Bond Strength

1303

Oral - 35

1304

INVESTIGATION OF POSSIBILITIES OF USING MOBILE BAGS IN DETERMINING DIGESTIBILITY IN DOG FOODS

1305

1306

MUSTAFA SELCUK ALATAS^a, OGUZHAN KAHRAMAN^a, FATMA INAL^a,

1307

NUTRETTIN GULSEN^a

1308

1309

^aselcuk university, faculty of veterinary, department of animal nutrition

1310

1311

selcukalatas@gmail.com

1312

This study was **conducted** to determine dry matter digestibilities of dog foods by in vivo and mobile nylon bag methods and compare them. **Thirty** of castrated young mature male dogs at approximately 15-30 kg live weights have been used. 6 dog foods have been evaluated. **In the study**, 30 young adult neutered male dogs with live weights ranging from 15-30 kg were used. A total of 6 dry dog foods, one of which is in pellet form, were examined. **The dry matter** digestibility rates of the dog foods determined by classical fecal collection method were 84.20, 81.17, 83.66, 83.47, 86.10 and 83.85%. The samples were fed to approximately 35 mg/cm² of dogs at 2x2 cm size mobile nylon bags with 35 μ pores and the bags were collected from the stool. The bags were washed for 4 minutes and 3 times changing water. Washing losses were also determined in the same way. The corrected for washing losses the dry matter digestibilities were determined as 83.06, 91.24, 79.53, 86.62, 94.58 and 87.21% (P<0.000), respectively. **The mobile nylon bag method** was used in dogs for the first time and it needs to be reconsidered and developed in terms of pore size of the bag, the amount of dog food put in it and the washing method.

1328

1329

Keywords: Digestibility, Dog Food, Mobile Bag

1330

1331

Oral - 36

1332

THE EFFECT OF HEAT TREATMENT APPLIED TO CEREALS USED IN DOG FOODS ON GELATINIZATION AND DIGESTIBILITY OF STARCH

1333

1334

FATMA INAL^a, ABDULLAH ÖZBILGIN^a, MUSTAFA SELÇUK ALATAŞ^a,

1335

OGUZHAN KAHRAMAN^a

1336

1337

^aselçuk üniversitesi veteriner fak. hay. bes. ve besl. hastalıkları a.b.d

1338

1339

abdullahozbilgin@gmail.com

1340

Barley, wheat, corn, rice, sorghum and oat were used in this study to determine the effects of soaking with hot water, cooking in water and extrusion processes on the gelatinization and digestibility of starch applied to cereals are used in dog food. **After grinding**, the grains were soaked with boiling water, cooked with water for 10 min and 20 min, and cooked with an extruder. The heat treated cereals were dried and ground. Dry matter, starch, gelatinized starch and in vitro starch digestibility analyzes were performed in all samples. **The amount of starch** in the heat treated cereals decreased. It was found that the richest cereal in terms of starch is rice and the poorest one is oat. The starch damage was determined as 4.64% in raw cereals, 30.99-31.83% in cereals cooked with water and 31.59% in extruded cereals(P<0.05). Heat treatments increased starch gelatinization by 581%. The highest gelatinization occurred in the oat. In vitro starch digestibility was found 14.62 in raw grains, 55.46 after cooking for 10 min and 72.47 mg maltose/mg starch after extrusion(P <0.05). Heat treatments increased starch digestibility by 308%. The highest starch digestibility occurs in the oat and the lowest in rice. **There was not** any gelatinized starch ratio difference between the cooking and extrusion, but starch digestibility was found higher after the extrusion process. Soaking with hot water was not effective in increasing starch gelatinization and digestibility.

1359

1360

Keywords: Dog Food, Cereals, Starch, Digestibility

1361

1362

Oral - 37

1363

A COMPARATIVE INVESTIGATION OF THE EFFECTS OF VARIOUS ANTIEPILEPTIC AGENTS IN THE RETINA AND CORNEAL LAYERS

1364

1365

AYŞE TUÇ YÜCEL^a, SEREN GÜLŞEN GÜRGEN^a, ALI CANSU^b

1366

1367

^aManisa Celal Bayar university

1368

^bKaradeniz technical university, Trabzon

1369

1370

ayse.tuc@gmail.com

1371

Aim: The number of studies suggesting that several antiepileptic agents used in the treatment of epilepsy may cause vision impairment has increased in recent years. We planned to perform a comparative investigation of the damage that may be caused in the retina and corneal layers by the antiepileptic agents sodium valproate (VPA), oxcarbazepine (OXC), levetiracetam (LEV) and topiramate (TPM) using nitric oxide synthase molecules. **Materials and Methods:** Five groups, each consisting of seven Wistar albino rats aged 21-24 days, were established. Group 1 represented the control group. Group 2 received VPA (300 mg/kg/day), Group 3 received OXC (30 mg/kg/day), Group 4 received LEV (50 mg/kg/day) and Group 5 received TPM (100 mg/kg/day) in two doses by gavage. At the end of the study period the eye tissues were removed and stained immunohistochemically with inducible nitric oxide synthase (iNOS) and endothelial nitric oxide synthase (eNOS) antibodies. **Results:** In contrast to the control group, quite powerful iNOS immunoreaction was observed in the inner plexiform layer of the retina in the VPA, OXC and LEV groups. Moderate reaction was observed in the VPA and OXC groups and weak reaction in the LEV group in the rod and cone layer. Moderate iNOS expression was observed in the inner and outer plexiform layers of the retina, and widespread weak iNOS expression in the rod and cone layers in the TPM group. Moderate release was observed at eNOS immunoreaction in the plexiform layer of the control group retinas, while weak reaction was observed in the outer plexiform and rod and cone layers. Moderate eNOS expression was noted in the inner plexiform layers in the VPA and OXC groups, while weak to moderate expression was observed in the inner plexiform layer in the TPM group.

1394

1395

Keywords: Epilepsy, Antiepileptic Agent, Sodium Valproate, Oxcarbazepine, Levetiracetam, Topiramate, Retina Corneal Layers

1396

1397

SULFASALAZINE MAY PREVENT DISSEMINATED INTERVASCULAR COAGULATION IN EXPERIMENTAL ENDOTOXEMIC RATS

EMRE BAHÇIVAN^a, HATICE ESER FAKI^a, BURAK DIK^a

^adepartment of pharmacology and toxicology, veterinary faculty, university of selcuk

emrebahcivan@selcuk.edu.tr

Disseminated intravascular coagulation is one of the most obvious causes of death in endotoxemic patients. The aim of the current study is to determine the effects of sulfasalazine treatment at different times on hematological, biochemical and coagulation factors. The **study** was performed on 30 Wistar albino rats and they were divided into 4 groups Groups; healthy control (HC), endotoxemic control (LPS) that was administration LPS (4 mg / kg, single dose), prophylactic sulfasalazine treatment group (SL5) that was treatment sulfasalazine (300 mg / kg, single dose daily) for 5 days before the endotoxemia model (4 mg / kg, single dose LPS) and sulfasalazine treatment group (LS) that was treatment sulfasalazine (300 mg / kg, single dose) at the same time with the endotoxemia model (4 mg / kg, single dose LPS). The blood was collected at 3 and 6 hours after the last applications in the all groups. Hematology, biochemical and coagulation [antithrombin (AT) III, fibrinogen and protein C] factors were measured in the blood samples. **The decreased** in fibrinogen level was partially inhibited in LS and SL5 groups. In addition, ATIII level in SL5 group was similar HC group at 3 hours. Protein C levels of SL5 group at 3 hours and LS group at 6 hours, was close to HC group. Sulfasalazine treatment partially elevated liver enzymes and the endotoxemia reduced erythrocyte, leukocyte and thrombocyte levels but decreased leucocyte levels were inhibited by sulfasalazine application (LS and SL5) at different times. all biochemical values were within the reference range reported for rats. **In conclusion**, application of sulfasalazine at different times may prevent acute intravascular coagulation on endotoxemia. In addition, the use of sulfasalazine at different times and at different doses may reduce the morbidity and mortality of the disease.

Keywords: Endotoxemia, Sulfasalazine, Hematology, Biochemistry, Coagulation

SOME REPRODUCTIVE PARAMETERS IN PRIMIPAROUS AWASSI SHEEP

ŞEREF İNAL^a, BEHIÇ ÇOŞKUN^b, DURSUN ALI DİNÇ^c, MUSTAFA ÇAM^a

^aselçuk university veterinary faculty animal science department

^bselçuk university veterinary faculty animal nutrition department

^cselçuk university veterinary faculty gynecology department

mustafa0609@hotmail.com

In this study, it was investigated reproductive parameters, birth weight, weaning weight and survival rates of 985 primiparous Awassi sheep, done by controlled mating, which were transferred from Ceylanpınar Agricultural Enterprise to İlci Çiçekdağı Agricultural Enterprise where the climate is colder and more rainy than Ceylanpınar, arid and hot climate conditions, to check their adaptation and productivity in colder climates. **Estrous rate**, birth rate, lambs born per ewe were calculated as %96.45, %95.47, 1.08 respectively. Twinning rate was found as %8,27. It was determined that %52,79 of lambs from the ewes were male. **32 of 34** breeding rams (%94,12) used in this study were able to mated with the ewes. Average number of ewes mated to the ram was 29,7 and maximum number was 75. The fertility rate of rams were between %27,78 and %100. **Even though** stillbirths were %0.12 for single lambs and %4 for twin lambs; mortality rate of lambs until weaning were found as %22,74 for single lambs and %20.14 for twin lambs. **Lamb birth and weaning weights** were determined as 4,756 and 28,436 kg for single lambs; 3.822 and 27.740 kg for twin lambs; 4,469 and 27,740 kg for females; 4.754 and 28,861 kg for males. The differences between lambing type and sex groups for birth weight were found really important ($P < 0.000$); but not for lamb weaning weight ($P > 0.05$). **As a result**; it can be said that reproductive parameters of Awassi sheep raised in the unnormal climate conditions such as rainy and cold weathers were not affected as much as expected. Obtained satisfying results from the study regarding to reproductive parameters in different climates, it can be also concluded that genotypes and preservation of Awassi population in Ceylanpınar Agricultural Enterprise was really in good conditions.

Keywords: Awassi Sheep, Primiparous, Reproductive Performance, Birth Weight, Weaning Weight

1470

Oral - 40

1471

GREEN SYNTHESIS OF SILVER NANOPARTICLES USING THYMBRA SPICATA VAR. SPICATA L. LEAF EXTRACT AND EVALUATION OF THEIR BIOLOGICAL ACTIVITIES

1472

1473

1474

FATİH ERCİ^a, RABİA ÇAKIR KOÇ^b

1475

1476

^aDepartment of biotechnology, faculty of science, Necmettin Erbakan university

1477

^bDepartment of bioengineering, faculty of chemical and metallurgical engineering, Yıldız
technical university

1478

1479

1480

ferci@konya.edu.tr

1481

In recent years, the use of metal nanoparticles for industrial applications has become an important field of study. The use of plant extracts in the development of fast, clean, non-toxic and eco-friendly process for nanoparticle synthesis has gained great importance. Here, for the first time, silver (Ag) nanoparticles (NPs) were green synthesized using Thymbra spicata leaf extract, also known as an important medicinal and aromatic plant species, is grown naturally in semi-arid climatic conditions of Turkey. Leaves and flowers are widely used for food, drink and medical purposes. Previous studies have reported antimicrobial analyses of essential oil, hydrosol and solvent-based extracts of T. spicata. The aqueous plant extract acted as reducing agent in the synthesis of nanoparticles. Characterization of the synthesized nanoparticles was performed by spectroscopic (UV-visible spectroscopy, FT-IR spectroscopy and XRD) and microscopic (AFM, SEM and TEM) analysis. Also, the average size, size distribution and PDI of the synthesized AgNPs in reaction mixtures were investigated by dynamic light scattering (DLS). The morphological characterization and size variation of the synthesized AgNPs with change in the extract quantity was carried out by microscopic analysis using TEM. The results showed that T. spicata leaf extracts mediated nanoparticle synthesis was cheap and reliable in room temperature conditions without any requirement for highly toxic chemicals. The synthesized AgNPs were studied for their antibacterial activities towards Gram-positive and Gram-negative bacteria. Also, we have carried out a comparative study on the cytotoxic effects of the synthesized AgNPs. The present study **demonstrates** that the size and shape are important parameters in controlling biological activities of nanoparticles.

1482

1483

1484

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

Keywords: Thymbra Spicata, Green Synthesis, Silver Nanoparticles, Biological Activity.

1506

1507

1508

Oral - 41

1509

MEASURING NITRATE CONSUMPTION BY *C. VULGARIS* USING NO₃- ION SELECTIVE ELECTRODE (ISE)

1510

1511

VILDAN ERCI^a, DIDEM ÖZÇİMEN^b

1512

^aDept. of soil science and nutrition, faculty of agriculture, Selcuk university

^bDept. of bioengineering, faculty of chemical and metallurgical engineering, Yıldız technical
university

1515

1516

1517

vildanerci@selcuk.edu.tr

1518

1519

1520

1521

1522

1523

1524

1525

1526

1527

1528

1529

1530

Microalgae can use substances such as nitrate in wastewater as nutrient and can continue both to purify the wastewater and their development. The reusability of wastewater makes its importance undeniable when considering the increasing population and living conditions. In this study, NO₃- ion selective electrode (ISE) that is used for selective determination of nitrate changes induced by *C. vulgaris* microalgae in a solar bioreactor containing 70 liters growth medium. Nitrate consumption by *C. vulgaris* microalgae in the solar bioreactor (designed specially and increased effectiveness) after approximately 60 days of ISE measurements revealed that the NO₃- concentration in the solar bioreactor decreased from 5 mg/L to 3 mg/L. **As a result**, it can be suggested that integration of nitrate ISE electrode into the solar bioreactor developed here would be useful for real time follow-up nitrate consumption at very close and certain intervals, and ISEs such as phosphate and pH should also be tested and integrated into the system.

1531

Keywords: Microalgae, *C. Vulgaris*, Solar Bioreactor, NO₃- ISE.

1533

1534

Oral - 42

1535

**MEMORY-ENHANCING, ANXIOLYTIC AND ANTIDEPRESSANT
EFFECTS OF INHALED ACHILLEA PSEUDOALEPPICA ESSENTIAL OIL
ON SCOPOLAMINE-INDUCED AMNESIC RATS**

1536

1537

1538

EMEL AKBABA, EYUP BAGCI

1539

1540

^aFırat Üniversitesi

1541

1542

emel19aydin@gmail.com

1543

1544

1545

1546

1547

1548

1549

1550

1551

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562

1563

1564

1565

1566

Achillea pseudoaleppica (Asteraceae) is an endemic plant species of Turkey. Several species of the genus has been used in folk medicine in the treatment of wounds, bleedings, headache, inflammation, pains, spasmodic diseases, flatulence and dyspepsia. In the present study, memory-enhancing, anxiolytic and antidepressant-like activities of *A. pseudoaleppica* essential oil on scopolamine-induced amnesic rats are assessed. In order to evaluate memory- enhancing activities of *A. pseudoaleppica* essential oil (1% and 3%, for 21 continuous days) Y-maze and radial arm maze tasks were used. As expected, scopolamine administration decreased spontaneous alternation percentage in Y-maze test and increased the number of working memory errors and the number of reference memory errors in radial arm-maze test. *A. pseudoaleppica* essential oil used in this study displayed an improved effect on acquisition of the short-term memory of scopolamine-treated rats within the Y-maze task as evidenced by increased spontaneous alternation percentage. *A. pseudoaleppica* essential oil administered rats also showed significant improvement of reference memory formation, but not working memory formation in radial arm maze test. Furthermore, *A. pseudoaleppica* essential oil induced anxiolytic effect on elevated plus maze test as evidenced by increased percentage of open arm time, open arm entries and the number of crossings as compared to scopolamine-alone treated rats. *A. pseudoaleppica* essential oil also exhibited antidepressant-like effects as evidenced by decreased immobility time on forced-swimming test. These results suggest the neuroprotective potential of *Achillea pseudoaleppica* essential oil on scopolamine-induced amnesia in rats and could act as a promising therapeutic agent for the neurological abnormalities related to Alzheimer's disease.

1567

1568

Keywords: *Achillea Pseudoaleppica* Essential Oil, Spatial Memory, Alzheimer's Disease, Anxiety, Depression

1569

1570

1571

1572

THE EFFECTS OF JUNIPERUS OXYCEDRUS SSP OXYCEDRUS ESSENTIAL OIL ON MEMORY, ANXIETY AND DEPRESSION IN AMNESIC RATS

1573

1574

1575

EMEL AKBABA^a, SAFEEN AHMED OTHMAN^b, EYUP BAGCI^c

1576

1577

^afirat üniversitesi

1578

1579

eydin@firat.edu.tr

1580

1581

1582

1583

1584

1585

1586

1587

1588

1589

1590

1591

1592

1593

1594

1595

1596

1597

1598

1599

1600

1601

1602

1603

In the present study, memory-enhancing, anxiolytic and antidepressant-like activities of Juniperus oxycedrus ssp oxycedrus (Cupressaceae) essential oil on scopolamine-induced amnesic rats are assessed. Scopolamine is a drug comprising highly amnesic properties; therefore it has been used to create dementia animal model. Scopolamine is known to induce memory impairment as well as anxiety and depressive-like behavior in animals and humans. In this study, we evaluated the neuroprotective effects of Juniperus oxycedrus ssp oxycedrus essential oil in scopolamine treated rats. Two different doses, 1% and 3%, of essential oil were administrated to the rats by inhalation for 21 continuous days prior to the behavioral analysis. Y-maze task and radial arm maze task were used to evaluate spatial memory performances of the rats. As expected, scopolamine treated rats exhibited the following: decreases in spontaneous alternation percentage in Y-maze task, and increases in working memory errors and reference memory errors in radial arm maze task. Juniperus oxycedrus ssp oxycedrus essential oil treated rats exposed to scopolamine showed significant improvement in spatial memory function as compared to scopolamine-alone treated rats. Furthermore, anxiety and depressive like symptoms were assessed by elevated plus maze task and forced swimming task. Juniperus oxycedrus ssp oxycedrus essential oil inhalation significantly increased the number of open arm entries in elevated plus maze, indicating anxiolytic behavior. The essential oil also reduced the immobility time and increased the swimming time in forced swimming test indicating antidepressant activity. **Taken together**, Juniperus oxycedrus ssp oxycedrus essential oil could be a good candidate for the treatment of dementia and deserves further investigation.

1604

1605

1606

Keywords: Juniperus Oxycedrus Ssp Oxycedrus Essential Oil, Memory, Dementia, Anxiety, Depression

1607

SHOOT REGENERATION OF HYPERICUM PERFORATUM L. FROM AXILLARY BUDS

ŞEYMA ÖNLÜ^a, HUSSEIN ABDULLAH AHMED AHMED^b, SERKAN URANBEY^c

^aScience and art faculty, department of molecular biology and genetics, Muş Alparslan
university, Muş, TURKEY

^bDept. of field crops, faculty of agriculture and natural sciences, Uşak university, Uşak,
TURKEY

^cDepartment of field crops, faculty of agriculture, Ankara university, Ankara, TURKEY

s.sofuoglu@alparslan.edu.tr

Hypericum perforatum L. is a medicinal plant, spread out in Turkey. Hypericum species have medicinal properties such as antidepressant, antibacterial, antiviral and has been used traditionally for many years in the world. Hypericum perforatum L. is a traditional medicinal plant containing biologically active and numerous secondary metabolites with anti-cancer, anti-depressive and anti-viral properties. The goal of the research was also to achieve high compact calli production and regeneration of *H. perforatum L.* by somatic embryogenesis or organogenesis. **Material-Methods:** The seeds of *H. perforatum L.* were collected from Uşak University Campus in Turkey. The seeds were treated with 1.5 g / L GA3 solution in 100 ml of distilled water for 2 days and then sterilized in 70% bleach for 20 minutes. The sterilized seeds were transferred and germinated on LS medium containing 3 % sucrose and 0.6 % agar. Axillary buds explants were isolated from germinated seeds for callus formation and cultured on LS medium supplemented with different concentrations of BAP (0.5 mg / L, 1 mg / L and 2 mg / L) and 0.1 mg/L NAA. All explants were kept at growth room with photoperiod for 16 hours light and 8 hours dark, at 24±2 °C. **Results and Discussion:** The best callus formation was obtained on LS medium containing 1 mg/L BAP and 0.1 mg/L NAA. The best adventitious shoot regeneration was also achieved on LS medium with 1mg/L BAP. The shoots derived from regenerative calli were rooted on MS medium supplemented with 1.5 mg / L IBA, and the roots of the plants were provided 8 days after culture initiation. The rooted plants were acclimatized adaptation in a mixture of peat, perlite and field soil (in a 1:1:1 ratio) in the growth chamber with high survival ratio (100%).

Keywords: Hypericum Perforatum L., Medicinal Plant, In Vitro Culture, Shoot Regeneration

ETHANOL INDUCED CHANGES IN THE EXPRESSION OF SELECTED GENES DOWNSTREAM TO NRF2 PATHWAY IN GRANULOSA CELLS IN VITRO

MD. MAHMODUL HASAN SOHEL, MUSTAFA ABDULKAREEM SALMAN,
ABDURRAHMAN AYVAZ

Erciyes university

sohel.mmh@gmail.com

Alcoholic beverages are a part of the social life in the west. However, chronic alcoholism may cause severe mental and physical damages including reproductive disorders. Granulosa cells (GCs) are the major cellular component in a follicular microenvironment and played an indispensable role in the maturation and protection of oocyte, the release of the oocyte, and the formation of corpus luteum. Therefore, the health and function of GCs dictate the outcome of a reproductive cycle as well as the pregnancy. This study was conducted to investigate the effects of chronic alcohol consumption on the function of ovarian GCs. For this purpose, we used in vitro culture of primary GCs from the bovine origin and different concentration of ethanol (0, 50, 200, and 500) to simulate the effects of alcohol consumption. Subsequently, cell viability and proliferation, mitochondrial activity, and expression of candidate genes were investigated. The results showed that there was a gradual loss of cell viability with the increase of the ethanol concentration, i.e. lowest viability was observed at the highest concentration (1000 mM), which is further supported by cell proliferation assay. Mitochondrial activity decreased significantly at higher concentrations. Candidate gene expression measured by qRT-PCR revealed that expression of NRF2 decreased significantly ($p < 0.05$) in ethanol exposed cells compared to the cells in control group whereas the expression of KEAP1, negative regulator of NRF2, increased significantly ($p < 0.05$) at 50 and 200 mM, however, no significant difference was observed at 500 mM concentration. The expression of antioxidant genes, downstream to Nrf2-pathway activation, showed that overall expression was decreased at 200 and 500 mM, however, increased at 50 mM concentration compared to the control. **The result of this study** provoked us to postulate that ethanol exposure decreases the ability of GCs to handle stress by downregulating the expression of genes involved in Nrf2-pathway.

Keywords: Alcohol, Nrf2-Pathway, Oxidative Stress, Granulosa Cells

Acknowledgements: This study was supported by the Erciyes University Scientific Research Projects Coordination Unit, Project No: 6734

1686

Oral - 46

1687

TIME AND DOSE-DEPENDENT EFFECTS OF OXIDATIVE STRESS INDUCED BY HYDROGEN PEROXIDE IN OVARIAN GRANULOSA CELLS

1688

1689

1690

MD. MAHMODUL HASAN SOHEL, YUSUF KONCA, MEHMET ULAS CINAR

1691

1692

Erciyes university

1693

1694

sohel.mmh@gmail.com

1695

1696

1697

1698

1699

1700

1701

1702

1703

1704

1705

1706

1707

1708

1709

1710

1711

1712

1713

1714

1715

1716

1717

1718

1719

1720

When the production of prooxidant molecules such as reactive oxygen species (ROS) overwhelmingly exceeding the cellular antioxidant capability, it potentially leads to a condition known as oxidative stress. Physiological level of ROS plays important roles in the reproductive process including follicular development, granulosa cell expansion, oocyte maturation, ovulation, and pregnancy. However, consistent production of an elevated level of ROS may disrupt the cellular homeostasis and results in oxidative stress. Many chemical compounds have been shown to be a potential inducer of oxidative stress in mammalian cells by increasing ROS level including hydrogen peroxide H₂O₂. Micromole of H₂O₂ is able to generate excessive ROS that may cause peroxidative damage to the cellular macromolecule and results apoptosis. The oxidative effects of H₂O₂ have been investigated in different types of cells, however, time and dose-dependent effects of oxidative stress induced by H₂O₂ in granulosa cells (GCs) have not been investigated yet. GCs were exposed to different concentrations of H₂O₂ (0, 100, 200, and 500 μM) for different time points (4 h, 12 h, and 24 h) and viability, cytotoxicity, and ROS accumulation was measured using trypan blue, WST-1 kit, and H₂DCFDA, respectively. The results showed that there were time and dose-dependent loss of cell viability in GCs due to H₂O₂ exposure. For instance, lower cell viability was observed at 500 μM concentration and 24 h time point while highest viability was observed at 100 μM concentration. Similar results were observed in cytotoxicity experiment. Analysis of ROS accumulation showed a concentration-dependent increase, i.e. higher ROS at higher concentration, and time-dependent decrease, i.e. lower ROS at higher time points, of ROS accumulation in GCs. **Our results** confirm that higher concentration fo H₂O₂ is highly cytotoxic and cause a loss of cell viability of GCs by generating excessive ROS in a time and dose-dependent manner.

1721

1722

Keywords: Oxidative Stress, Cell Viability, Granulosa Cells

1723

THE EFFECTS OF CHRYSINE ON CYPERMETHRIN-INDUCED ACUTE TOXICATION IN RAINBOW TROUT (ONCORHYNCHUS MYKISS)

ÖZGE CERİT^a, FERIDE KOC^b

^aDept. of pharmacology and toxicology, faculty of veterinary medicine, Erciyes university

^bDept. of pharmacology and toxicology, faculty of veterinary medicine, Erciyes university

feridekoc@yahoo.com

Cypermethrin (SP) is a pyrethroid insecticide. It is high toxic for fishes and aquatic live, also. Chrysine (KR), anti-inflammatory, antiviral, anticarcinogen and antioxidant characteristic, is a flavonoid and it is obtained from propolis, some mushroom, and some natural plants. The purpose of this study is to determine effects of the KR on SP-induced acute toxication in fish. **In study**, 60 fish divided into 6 groups (n=10). Group 1: control, Grup 2: 0.4 µg/L SP; Grup 3: 50 mg/kg KR; Grup 4: 0.4 µg/L SP+50 mg/kg KR; Grup 5: 0.4 µg/L SP+100 mg/kg KR; Grup 6: 100 mg/kg KR. After, blood and tissues was collected. It was determined cytokines (IL-1β, IL-6, IL-10 and TNF-α), ALT, AST in serum and MDA, SOD, GSH and CAT levels in tissues (liver and kidney). **In this study**, AST, ALT, cytokines (except IL-10) and MDA increased, SOD, GSH and CAT decreased in SP group. On the contrary, when we compared KR groups with SP group, KR significantly decreased IL-1β, IL-6, and TNF-α, ALT, AST and MDA. Also, GSH, SOD and CAT increased depending on dose in KR groups. **In conclusion**, KR can prevent tissue damage via antiinflammatory and antioxidative effects in SP-induced acute toxication in fish.

Keywords: Mice, Anethum Graveolens, Paracetamol, Lipid Peroxidation, Toxication. Fish, Cypermethrin, Chrysine, Antioxidant, Cytokine.

Acknowledgements: This study is the MSc thesis of Özge CERİT. This study was supported by the Scientific Research Council of Erciyes University, K

THE PROTECTIVE EFFECTS OF ANETHUM GRAVEOLENS ON PARACETAMOL-INDUCED ACUTE TOXICATION IN MICE

SERAP KORKMAZ^a, FERIDE KOC^b

^aSafiye Çikrikçioğlu vocational collage, department of veterinary, Erciyes university

^bDept. of pharmacology and toxicology, faculty of veterinary medicine, Erciyes university

feridekoc@yahoo.com

Objective: Parasetamol (PT), paraaminophenol group is a drug and cause damage on liver tissue when used in high doses with formation of its toxic metabolite n-asetil-p-benzokuinonenimin. Anethum graveolens (AG), antiinflammatory and antioxidant characteristic, is a plant. The purpose of this study is to determine effects of the AG on PT-induced acute toxication in mice. **Methods:** In study, 50 mice divided into 5 groups (n=10). Group I: control, Group II: 400 mg / kg a single dose PT orally, Group III: PT + AG1 (100 mg / kg AG), Group IV: PT + AG2 (200 mg / kg AG), Group V: AG2 (200 mg / kg). After, blood and tissues was collected. It was determined serum cytokine (IL-1 β , IL-6, IL-10 and TNF- α), ALT, AST and MDA, SOD, GSH in tissues with histopathological examinations. **Results:** AST, ALT, cytokines and MDA increased, SOD and GSH decreased in PT group. On the contrary, when we compared AG groups with PT group, AG significantly decreased cytokine, ALT and AST. The MDA decreased, and GSH, SOD increased depending on dose. Histopathological findings are consistent with biochemical findings. **Conclusion:** AG can prevent organ damage via affected cytokine answer and oxidative stress in PT-induced acute toxication in mice.

Keywords: Mice, Anethum Graveolens, Paracetamol, Lipid Peroxidation, Toxication

Acknowledgements: This study is the MSc thesis of Serap KORKMAZ. This study was supported by the Scientific Research Council of Erciyes University

1785

Oral - 49

1786

ANTIMICROBIAL APPLICATION OF GREEN SYNTHESIS OF ZINC OXIDE NANOPARTICLES BY EUPHORBIA MACROCLADA EXTRACT

1787

1788

SUZAN SAHIN DOGAN^a, AYTAC KOCABAS

1789

1790

Department of biology, karamanoğlu mehmetbey university, Karaman, TURKEY

1791

1792

sdogan@kmu.edu.tr

1793

1794

1795

1796

1797

1798

1799

1800

1801

1802

1803

1804

1805

1806

1807

1808

The purpose of the study is to investigate antimicrobial activity of biologically synthesized Zinc Oxide (ZnO) nanoparticles. ZnO nanoparticles have been of great interest due to their unique properties such as chemical stability, virtuous conductivity, catalytic and antibacterial, antifungal anti-inflammatory activities. ZnO nanoparticles are used as antimicrobial agent in various areas like medical, pharmaceutical, food and agricultural science. Green nanotechnology provides easy shape and size control of synthesized nanoparticles in addition to eco friendly synthesis of nanoparticles by utilizing biological materials such as microorganisms, fungus and plants. The aqueous extract of Euphorbia macroclada was applied as biological capping and surface stabilizing agent for the synthesis of ZnO nanoparticles. Soxhlet extractor was performed to acquire the aqueous extract of Euphorbia macroclada. Creamish-white colored precipitation was confirmed the synthesis of nanoparticles. UV-vis spectrophotometer, Zeta sizer, FTIR and XRD were used ZnO nanoparticles characterization. **Antimicrobial activities** of ZnO nanoparticles were investigated on Staphylococcus aureus, Bacillus subtilis, Pseudomonas aeruginosa and Escherichia coli by using well diffusion method.

1809

1810

Keywords: Zinc Oxide, Nanoparticle, Euphorbia Macroclada, Antimicrobial

1811

THE EFFECTS OF PROPHYLACTIC AND THERAPEUTIC**SULFASALAZINE IN LIVER OF EXPERIMENTAL ENDOTOXEMIC RATS**

HATICE ESER FAKI^a, GONCA SÖNMEZ^b, EMRE BAHÇIVAN^c, BURAK DİK^d^aSelcuk university veterinary facultyhaticeeser@selcuk.edu.tr

Lipopolysaccharide (LPS) is obtained from gram (-) bacteria cell wall including E. coli, and it has an important role in the formation of septic shock and endotoxemia. In this study, it was aimed that determination of prophylactic and therapeutic effects of sulfasalazine in the experimentally induced endotoxemia in rats. **The study** was carried out with 4 groups of 30 Wistar albino rats. Groups was established as control group (C), experimental endotoxemia model group (EE) with LPS (4 mg/kg, single dose), prophylactic sulfasalazine treatment in experimental endotoxemia model group (PSE) (300 mg/kg, single dose daily, for 5 days before the experimental endotoxemia model) and lastly sulfasalazine treatment in experimental endotoxemia model group (SE) (300 mg/kg, single dose daily, at the same time as endotoxemia model). The rats were euthanized at 6 hours after the last applications in the all groups, and mRNA levels of IκB, EDN1, BCL2 and BAX gene expression were examined from liver tissues by real-time PCR. **It was determined** that the level of IκB in the liver increased with endotoxemia and this increase was inhibited by the application of prophylactic sulfasalazine. Expression of EDN1, which is important for the perfusion of the organs, statistically changed by prophylactic and therapeutic sulfasalazine applications. Anti-apoptotic BCL-2 increased in liver mRNA expressions in SE group and BAX expression partially decreased in the PSE and SE groups. **As a result**, prophylactic and therapeutic sulfasalazine may prevent apoptosis and inflammation via inhibiting IκB in endotoxemia. The effects of different doses and durations of sulfasalazine should be determined on other organs in the endotoxemia model.

Keywords: Endotoxemia, Sulfasalazine, Ikb, BCL-2, Bax

1845

Oral - 51

1846

PHARMACOKINETICS OF CEFTRIAZONE FOLLOWING SINGLE ASCENDING INTRAVENOUS ADMINISTRATION IN SHEEP

1847

1848

DUYGU DURNA ÇORUM^a, ORHAN ÇORUM^b, HATİCE ESER FAKI^c,

1849

EMRE BAHÇIVAN^c, AYŞE ER^c, KAMİL ÜNEY^c

1850

^aAraç Rafet Vergili vocational high school, university of kastamonu, Kastamonu, TURKEY

^bDept. of pharmacology and toxicology, faculty of veterinary medicine, university of

Kastamonu, Kastamonu, TURKEY

^cDept. of pharmacology and toxicology, faculty of veterinary medicine, university of Selcuk,

Konya, TURKEY

1856

haticeeser@selcuk.edu.tr

1857

1858 **Ceftriaxone** (CTX) is a third-generation cephalosporin, which used more frequently

1859 in veterinary clinical practice because of the activity of its broad spectrum

1860 including Gram-positive and Gram-negative microorganisms and the low toxicity.

1861 Although different dose levels of CTX may be used in the treatment of infections

1862 caused susceptible pathogens in sheep, the pharmacokinetics of ascending doses of

1863 CTX in sheep was not found in literature research. The objective of this study was

1864 to evaluate the pharmacokinetics of CTX following intravenous administrations of

1865 single ascending doses in sheep. In this study, six clinically healthy Akkaraman

1866 sheep (2.4±0.4 years and 50±3 kg body weight) were used. CTX was administered

1867 intravenously to each sheep at doses of 20, 40 and 80 mg/kg in a crossover design

1868 with a 15-day washout period. Plasma concentrations of CTX were measured using

1869 the HPLC-UV method. Pharmacokinetic parameters were calculated by non-

1870 compartmental analysis. CTX is well tolerated following administration at doses of

1871 20, 40 and 80 mg/kg. While mean residence time (MRT) of CTX at the dose of 80

1872 mg/kg was longer than those in other doses (P<0.05), its elimination half-life

1873 (t_{1/2λz}) following the administration at doses of 40 and 80 mg/kg were longer than

1874 that in 20 mg/kg dose (P<0.05). While total clearance of CTX after 40 and 80

1875 mg/kg doses decreased significantly compared to 20 mg/kg dose, its dose-

1876 normalized area under the plasma concentration-time curve (AUC_{0-∞}) at 80 mg/kg

1877 dose increased significantly compared to 20 mg/kg dose. The volume of

1878 distribution at steady-state of CTX was not significantly different between the dose

1879 groups (P >0.05). Our study showed that CTX can be used at 12 h interval to

1880 maintain T>MIC of >40 for the treatment of infections caused bacteria with MIC

1881 values ≤2, ≤4, and ≤16 µg/mL, respectively, after 20, 40, and 80 mg/kg doses of

1882 CTX. **This information** may be helpful in adjusting the dosage regimen but there is

1883 a need for future work.

1884

1885 **Keywords:** Ceftriaxone, Ascending Doses, Pharmacokinetics, Sheep

1886

Oral - 52

1887

DOXORUBICIN LOADED CLAMSHALL TYPE FOUR DITHIOL-O-CARBORANE BRIDGED METALLOPHthalOCYANINES FOR PHOTODYNAMIC THERAPY

1888

1889

1890

BERKE BILGENUR ŞENER^a, SEVIL ŞENER^b, ABDULLAH TAHIR BAYRAÇ^a

1891

^a Department of bioengineering, faculty of engineering, Karamanoğlu Mehmetbey university

1893

^b Community college of aliaga, Ege university

1894

1895

bilgenurk@gmail.com

1896

1897

1898

1899

1900

1901

1902

1903

1904

1905

1906

1907

1908

1909

1910

1911

1912

1913

1914

1915

1916

1917

1918

INTRODUCTION: Cancer is one of the leading causes of death worldwide. Among cancer therapies, photodynamic therapy (PDT) was shown to be a clinically approved minimally invasive therapeutic method which can show a selective cytotoxic activity toward malignant cells. Phthalocyanines are used as photosensitizers which are components of PDT and with light and oxygen they initiate a photochemical reaction that culminates in the generation of a highly reactive product termed singlet oxygen (1O_2). **METHODS:** In this study clamshell-type 1,2-bis(2-thiol)-o-carborane bridged binuclear metallophthalocyanines (M = Zn, Co; Zn-Pc and Co-Pc, respectively) were synthesized. Structures of the target compounds were characterized by UV-Vis, FT-IR, MALDI-TOF mass and H-NMR spectroscopy techniques. Doxorubicin (DOX) loading on the Pcs were analyzed using spectrofluorometric techniques. Cytotoxicities of Pcs alone and Pcs loaded with DOX were tested using cell viability tests with N2a, MCF 7, HEK 293 and MIA PaCa-2 cells. All samples were prepared in two separate groups and one of the groups were subjected to PDT throughout the incubation (PDT (+) cells). **RESULTS:** Fluorescence intensity of DOX decreased by 12 % and 62 % for 1:1 and 1:10 DOX: Zn-Pc (mol: mol) combinations, respectively, and 25 % and 79 % in presence of Co-Pc (DOX: Co-Pc, 1:1 and 1:10, respectively). Viabilities of PDT (+) cells incubated with DOX loaded Pcs were lower when compared to PDT (-) cells. The most dramatic decrease was seen in MIA PaCa-2 cell line with Co-Pc PDT (+) with 54 % when compared to untreated cells. **CONCLUSION:** Zn-Pc and Co-Pc release DOX in response to light irradiation and this system could be a potential component of a stimuli responsive drug delivery system.

1919

1920

Keywords: Phthalocyanine, Photodynamic Therapy, Doxorubicin, Cancer Therapy

1921

In vitro maturation in mouse

MUSTAFA NUMAN BUCAK^a, ALİ ERDEM ÖZTÜRK^b

Selcuk University

mnumanbucak@gmail.com

As oocyte sources, we can use preantral or superovulated follicles. For using of preantral follicles, female mouse is killed by cervical dislocation at 14 days old. All selected follicles are placed into in vitro maturation conditions. Several effective maturation media are used for maturing of mouse oocytes. Follicles are cultured in an incubator for 12 days at 37°C, 100% humidity and 5% CO₂. After EGF and hCG are added the maturation medium from days 13, maturing follicles start to ovulate (14 hours later) and matured MII oocytes are released. For using of superovulated follicles; after hormonal treatments, ovaries are cleaned from fat and oviduct tissues. Then oocytes are removed with the help of 26G needles into washing solution. They are putted into 50µL culture medium and evaluated the expansions of oocytes for IVF. In this study, we explain detail procedures of in vitro maturation of mouse. **In vitro maturation** in mouse is different from some other mammals. Understanding of unknown mechanism in maturation stage requires effective studies with well controlled. In this way, mouse is getting more important, and can be used for developing new effective media and understanding mechanisms of embryo blocking/developing systems.

Keywords: Follicles, In vitro maturation, Mouse

1948

Oral - 54

1949

An investigation of blood iron, copper and zinc levels in sheep at region of Suruç

1950

1951

İBRAHİM HALİL ÇELİK^a, İLKER CAMKERTEN^b, GÜZİN CAMKERTEN^c

1952

TEKİN ŞAHİN^d

1953

^a Food, Agriculture and Animal Husbandry County principalship, Haliliye, Şanlıurfa

1954

^b **Supervisor**, Dept. of internal medicine, faculty of vet. med. ^c **Co-supervisor** Dept.

1955

Veterinary, Vocational School of Technical Sciences, Aksaray University, Aksaray,

1956

^d **Project Manager**, Dept. of internal medicine, faculty of vet. med, Siirt University, Siirt, Türkiye

1957

1958

1959

oguzalperen@hotmail.com

1960

1961

The case of **lack or excess of trace elements** in animals causes serious clinical disorders and significant economic losses. Imbalances of trace element in farm animals are results in all the animal productivity decline including hair loss, depigmentation, parakeratosis, continuous diarrhea, anemia, anorexia, bone formation disorders, tetany, the abortion which is not due to infection, fry efficiency as well as infertility, stunted and pica. **The project area** was identified as Suruc district of Sanliurfa. Aşağıoylum, Yönlü, Kaplan and Ayhan villages were selected as study area which belonging to Suruc. Randomly 25 blood samples were collected from the each village. **Serum iron, zinc and copper levels** of the four villages were measured as 67.37 ± 2.29 mg/dL, 77.29 ± 2.59 mg/dL and 157.64 ± 6.99 mg/dL, respectively. The levels of iron and copper were normal compared to the reference values, but the zinc levels were low. The copper levels of Aşağıoylum and Yönlü villages were significantly lower than those of the other regions ($P < 0.01$), and also zinc and iron levels of Kaplan region were significantly higher than those of the other regions ($P < 0.01$) in the analysis of variance of inter-regional. **As a result;** zinc levels were low and copper and iron levels were found to be normal in the sheep in Suruc. It was pointed out that the zinc supplementation in the area was priority important for the preventive medicine.

1979

1980

Keywords: Copper, Zinc, Iron, Sheep, Suruc, Sanliurfa.

1981

Acknowledgements: *This study is the MSc thesis of İbrahim Halil Çelik, DVM, MSc. This study was supported by the Scientific Research Council of Harran University with project number 13131.*

1982

1983

1984

1985

Investigation of iron, copper and zinc levels in sheep in district of Akçakale

METİN BOZ^a, İLKER CAMKERTEN^b

^aDoğa Veterinary Clinic, İstanbul & Health Science Institute, Harran University

^b**Supervisor**, Dept. of internal medicine, faculty of vet. med. Aksaray University

TÜRKİYE

ilkeracamkerten@gmail.com

Trace elements which constitute the basis of life have of great importance. Various diseases may occur in sheep and lambs in the absence of trace elements. Especially enzootic ataxia, swayback, alkali disease which occur in case of Cu, Zn and Fe deficiency, and anemia has of great importance in sheep breeding .Today's animal breeding, diseases caused by lack of trace elements has become as important as parasitic and infectious diseases. In this study, it is aimed to determine the blood serum iron, copper and zinc levels in healthy sheep in Akçakale region where this kind of investigation has not been done before. **Akçakale is district of Şanlıurfa** province and was chosen as a study area and the region was divided into three region. Total of 107 blood samples from the sheep, 34 samples from the first region, 37 samples from the second region and 36 samples from the third region, were taken. Analysis performed by ICP-OES, and serum iron, copper and zinc concentrations were measured in region 1, 2, and 3 were $109,44 \pm 20,40$ mg / dl, $58,55 \pm 13,15$ mg / dl and $59,60 \pm 8,38$ mg / dl, respectively. **When the results were evaluated**, serum iron, copper and zinc levels were determined lower compared to reference values. In variance analysis, serum zinc levels were determined to be lower (statistically significant, $p < 0,05$) compared to other regions. **As a result**, hidden deficiency in the terms of iron, copper and zinc in sheep can be considered, and especially zinc deficiency was more appealing in Akçakale region.

Keywords: Copper, Iron, Zinc, Sheep, Şanlıurfa, Akçakale

Acknowledgements: *This study is the MSc thesis of Metin Boz, DVM, MSc. This study was supported by the Scientific Research Council of Harran University with project number 12180.*

2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048

interim

POSTER PRESENTATIONS

2049

Poster - 1

2050

DNA LIBRARY DESIGN FOR SELEX USING BIOINFORMATIC TOOLS

2051

BERKE BILGENUR ŞENER^a, DENİZ YİĞİT^a, ABDULLAH TAHİR BAYRAÇ^a

2052

^aDepartment of bioengineering, faculty of engineering, Karamanoğlu Mehmetbey university

2054

2055

bilgenurk@gmail.com

2056

INTRODUCTION: Systematic evolution of ligands by exponential enrichment (SELEX) is used to select DNA aptamers and uses PCR method for enrichment of sequences that capture targets with their 3D structure. A 80 bases long sequence of DNA library 5'- F-Primer - (N45) – R-Primer-3', composed of two primer binding regions at both ends and 40-45 base long random sequence in between was designed using bioinformatic tools in this study. Forward primer (F-primer) was tagged with FITC on 5' end and was in the form of 5'- /5Fluor (N18) -3'. PCR is a major part of aptamer selection and therefore it is important to use primers and a library set that have high PCR efficiency. Since random region in the designed libraries already causes unwanted PCR products primer regions had to be carefully designed. **METHOD:** The primers of DNA library were designed to have appropriate annealing temperatures and GC content higher than 50%. Possibilities of G-quadruplex formation that could result from high G content of the sequences were calculated using QGRS Mapper software (bioinformatics.ramapo.edu/QGRS/) and sequences containing any G-quadruplex structure were eliminated. Additionally, heterodimers between two primer sequences and their complementary sequences on DNA library, self dimers and hairpins for each of the primer sequences, and hairpins within DNA library sequence were avoided. The stabilities of possible undesired structures within designed sequences at both 4°C and 37°C were analysed using NUPACK Nucleic Acid Package software (www.nupack.org/) and IDT DNA Technologies-OligoAnalyzer tool (www.idtdna.com/calc/analyzer). **CONCLUSION:** Two DNA library sequences were selected from five candidate sequences. The selected DNA library sequences could be used in various SELEX procedures for DNA aptamer development.

2080

Keywords: SELEX, DNA Library Design, Nupack

2082

2083

Poster - 2

2084

ANALYSIS OF PROTEIN AND MRNA EXPRESSIONS OF CYP1A1 AND CYP2E1 ENZYMES IN LIVER, COLON AND PROSTATE CANCER CELL LINES TO STUDY DRUG AND CARCINOGEN METABOLISM

2085

2086

2087

EMRE EVIN^a, SERDAR KARAKURT^b, ORHAN ADALI^a, MERVE AKKULAK^c

2088

2089

^aDepartment of biology, faculty of science and arts, middle east technical university

2090

^bDepartment of biochemistry, faculty of science and arts, selçuk university

2091

^cDepartment of molecular biology and genetics, faculty of science and arts, middle east technical university

2092

2093

2094

emrevin@gmail.com

2095

2096

Most of the marketed drugs are metabolized by Cytochrome P450s (CYPs) known as phase I enzymes. A phase I enzyme, CYP1A1 catalyzes the conversion of many pro-carcinogens and hydroxylation of many endogenous substrates. CYP2E1 is also important in the metabolism of drugs, pre-toxins and pro-carcinogens. The present study was aimed to describe the best cell line model for studying phase I xenobiotic metabolizing CYP1A1 and CYP2E1 enzymes and possible effects of xenobiotics on these enzymes. In this study HT29 and SW620 (colon); HEPG2 and HUH7 (liver); PNT1A and PC3 (prostate) cell lines were chosen. Then, mRNA and protein expressions of the CYP1A1 and CYP2E1 enzymes in these cell lines were analyzed by qRT-PCR and Western blotting techniques, respectively. It has been found that CYP1A1 protein was expressed in all cell lines and relative protein expression is highest in the HT29 (100%). In addition, HEPG2 (0.63 fold) and HUH7 (1 fold) expressed highest CYP1A1 mRNA. CYP2E1 protein expression was found in all cell lines at relatively high levels except PC3. CYP2E1 mRNA expression was significantly higher in HUH7 (1 fold) and HT29 (0.59 fold). **According to the results**, there was no significant correlation between mRNA and protein expressions. Thus choosing the best cell line as model depends on the purpose of the research. For studying metabolism of a chemical by CYP1A1 and CYP2E1 or effect of a chemical on translational regulation of these enzymes, it is better to consider protein expression levels. However, if the aim is to study effect of a chemical on transcriptional regulation of these enzymes, it is better to choose a cell line that expressing highest mRNA of gene of interest. Considering both mRNA and protein expression levels together, the best model cell lines for studying CYP1A1 and CYP2E1 are HT29 and HUH7, respectively.

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

2108

2109

2110

2111

2112

2113

2114

2115

2116

2117

2118

2119

2120

Keywords: Xenobiotic Metabolism, CYP1A1, CYP2E1, Western Blot, Qrt-PCR, In Vitro

2121

2122

Poster - 3

2123

COMPARATIVE EVALUATION OF SMALL T GENE AND LARGE T GENE BASED REAL-TIME PCR METHODS FOR THE DETECTION OF THE BK AND JC VIRUSES IN CLINICAL SAMPLES

2124

2125

2126

UĞUR TÜZÜNER^a, MEHMET ÖZDEMİR^b, MURAT ŞEVİK^c, MAHMUT BAYKAN^b

2127

2128

^aMedeniyet university, göztepe training and research hospital, department of medical microbiology

2129

2130

^bDept.of medical microbiology, meram medical faculty, Necmettin Erbakan university

2131

^cKonya veterinary control institute, molecular microbiology department

2132

2133

mehmetozdem@yahoo.com

2134

Objective: The aim of our study is to compare and evaluate the results of real-time PCR methods targeting the small and large T gene regions of the viral genome, taking into account polymorphisms occurring in the viral genome of BK and JC viruses. **Method:** Urinary specimen of 82 patients over 18 years old who were sent with suspicion of BKV and JCV to Molecular Microbiology laboratory of Necmettin Erbakan University, Meram Medical Faculty were taken into study. The small t gene was investigated using a commercial kit (LightMix, Roche, USA) and real-time PCR method. Large T gene-specific nucleic acid probes were investigated using the optimized in-house real-time PCR method using primers and probes as described by Dumoulin and Hirsch. Sequence analysis of the VP1 region was also performed using the Sanger method in 19 samples that has different PCR results. **Results:** BKV positivity in 9 samples and JCV positivity in 61 samples by real-time PCR method specific to small t gene region; BKV positivity in 21 samples and JCV positivity in 67 samples were determined by real-time PCR method specific to the large T gene region. Statistically, there are two methods; There was a significant difference for BKV, no significant difference for JCV. Sequence analysis is accepted as the standard method and the in-house method we designed with primers and probes; Sensitivity for BKV was 100%, specificity was 81.3%, positive predictive value was 33.3%, negative predictive value was 100%, sensitivity for JCV was 98.5%, specificity was 100%, positive predictive value was 100% and negative predictive value was 93.3%. **Conclusion:** It was thought that polymorphisms in the different gene regions targeted were responsible for the different outcomes obtained at the end of our study. With this sensitivity and specificity, our method is a candidate for routine diagnosis.

2135

2136

2137

2138

2139

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

2155

2156

2157

2158

2159

Keywords: BK Virus, JC Virus, Real-Time PCR, Polymorphism

2160

Poster - 4

2161

EFFECT OF BLOOD HORMONE LEVELS AND BODY CONDITION SCORE IN SUCCESS OF ARTIFICIAL INSEMINATION IN COWS

2162

2163

İLKER YAVAŞ^a, İSHAK GÖKÇEK^b, ECE KOLDAŞ^c, TUĞBA KORKMAZ YAVAŞ^d

2164

2165

^aMustafa Kemal Üniversitesi veteriner fakültesi dölerme ve suni tohumlama a.d

2166

^bMustafa Kemal Üniversitesi veteriner fakültesi fizyoloji a.d

2167

^cMustafa Kemal Üniversitesi veteriner fakültesi doğum ve jinekoloji a.d

2168

^dHatay il tarım müdürlüğü gıda kontrol laboratuari

2169

2170

ishakgokcek@hotmail.com

2171

Fertility rate with artificial insemination obtained depends on various factors. One of these factors is the body condition score. In this study, the body condition scores determined during artificial insemination, follicle development, steroid hormone (estrogen) and the impact on the level of estrus symptoms and aimed to reveal the possible cause of infertility. **In this study**, cows with no obstacles impeding fertility and at least one birth without any problems have been used at the private farms in Hatay Region. **Body Condition Score:** The oestrus assessments of cows that have been identified have been made according to the following criteria: Uterian flow, edema of Vulva, Vagina's color, Duldung reflex and diameter of follicle. **Cows** that are determined to be at estrus were inseminated by recto-vaginal method. **Estrogen** levels were determined by immunological methods (ELISA). **As a result**, it has been observed that if the body condition score is less than 2.5 in artificial insemination, the amount of estrogen is decreased, the first insemination pregnancy rate may decrease from 10% to 18% and the birth-first insemination interval may be prolonged.

2186

2187

Keywords: Body Condition Score, Infertility, Hormone Levels, Artificial Insemination.

2188

2189

Acknowledgements: This research was financed by Research Fund of Mustafa Kemal University (Project Number:13882)

2190

2191

Poster - 5

2192

THE HEMOGLOBINOPATHY PREVALENCE OF THE PATIENTS WITH SUSPECTED HEMOGLOBINOPATHY WHO HAVE APPLIED TO AFYON KOCATEPE UNIVERSITY PRACTICE AND RESEARCH HOSPITAL

2193

2194

2195

AHMET KAHRAMAN^a, MUSTAFA ÖRMECİ^b, ZELİHA ÇELİK^b, MUHAMMED KAYA^b

2196

2197

^aafyon kocatepe university

2198

^bafyon kocatepe university, student in faculty of medicine

2199

2200

akahraman57@yahoo.com

2201

Objectives and aimed: Hemoglobinopathies are the most common autosomal recessive hematological diseases in the world. According to World Health Organization data, the carrier frequency is around 7% in the world and 300.000-500.000 children with hemoglobinopathy are born every year. In our country, the mean β thalassemia carrier is 2%, and the most common of hemoglobinopathies are the sickle cell anemia (HbS) carrier (% 0.6). It is estimated that there are a total of 1,300,000 carriers and about 4,000 thalassemia patients in Turkey. In our country; the carriage of thalassemia in Çukurova, Mediterranean coastal strip, Aegean and Marmara regions is more common than in other regions. The aim of our study is to determine the prevalence of hemoglobinopathy in suspected patients who come to Hospital of Afyon Kocatepe University and to shed light on the establishment of control programs that have an idea about the frequency of hemoglobinopathy in Afyonkarahisar and its region and to prevent the birth of sick children. **Materials and Methods:** In our study, HbA and HbA2 values of 285 suspect patients who came to Medical Biochemistry Laboratory of Afyon Kocatepe University hospital were examined retrospectively according to archival records. Results: HbA and HbA2 values were found to be healthy in 176 and thalassemia minor patients in 109 (HbA2>3.5%) of the 285 cases. Conclusions: As a result, it was determined that the proportion of carriers of β -thalassemia in the suspect patients who applied to Medical Biochemistry Laboratory of Afyon Kocatepe University hospital was much higher than the average of Turkey (2%) and for this reason, it was **concluded** that the future suspect patients should be made aware of the issue and that the screening of premarital hemoglobinopathy is mandatory.

2224

2225

Keywords: Thalasemi Prevalence, Afyon Region, Hba2

2226

2227

Poster - 6

2228

ANTIOXIDANT PROPERTIES OF DIETARY FLAVONOIDS

2229

AHMET KAHRAMAN

2230

2231

Afyonkocatepe university

2232

2233

akahraman57@yahoo.com

2234

Recent studies have demonstrated that reactive oxygen species (ROS) such as superoxide anion radical, hydroxyl radical and lipid peroxidation play an important role in the pathogenesis of various diseases such as cancer, inflammatory diseases, cardiovascular and cognitive disorders. Nevertheless, the studies in recent years also indicated that widely consumed some foods contain some compounds capable of protecting individuals from the development of cancer, cardiovascular disorders and chronic diseases. Those compounds are called dietary antioxidants and their protective effects attributed to their scavenging activity. Flavonoids are a large group of antioxidants naturally occur in vegetables, fruits, cereal, tea and red wine. **In this study**, we want to present recent knowledge about structure, function and sources of flavonoids.

2245

2246

Keywords: Flavonoids, Antioxidant Properties, Relationship Of Structure- Activity

2247

2248

Poster – 7

2249

INVESTIGATION OF IMMUNOLOGICAL INFERTILITY IN NON-FERTILIZED COWS

2250

2251

ILKER YAVAŞ^a, BESTAMI KEMAL GÜMÜŞAY^b, ISHAK GÖKÇEK^c

2252

2253

^aMustafa kemal üniversitesi

2254

^bGıda tarım ve hayvancılık bakanlığı

2255

^cMustafa kemal üniversitesi

2256

^e

2257

ishakgokcek@hotmail.com

2258

The most important productivity of animals is the productivity of the offspring. Reproduction infertility is one of the major problems of income reduction in livestock enterprises. Causes of infertility in cows are very diverse. These include hereditary, functional (anoestrus, subestrus, cysts), infectious diseases, care and feeding disorders. It is noteworthy that in some cows the cases of repeat breeder have been observed despite the many seeds made even in the absence of these reasons. **In this study**, it was aimed to investigate the incidence of immunological infertility by determining immunological infertility by looking at immunoglobulin levels in repeat breeder cows. **In the study**, cows who had never been pregnant at least 3 times insemination in the private farms and did not have any disorders preventing fertility in the rectal and ultrasonographic examinations were used. Antisperm antibodies were detected by ELISA test in blood serum and cervical mucus samples obtained after insemination. **As a result**, 14.33% of sperm antibodies (ASA) immunity against sperm were detected in the sera of cows that were inseminated at least 3 times but not conceived, without any disease tested in normal field conditions. These animals have the potential to become pregnant with appropriate treatment (corticosteroid).

2275

2276

Keywords: Loss Of Fertility, Immunological Infertility, Immune Globulin, Artificial Insemination

2277

2278

Acknowledgements: This research was financed by Research Fund of Mustafa Kemal University (Project Number:13881)

2279

2280

2281

Poster - 8

2282

L-LDH GENE KNOCK-OUT OF LACTOBACILLUS HELVETICUS OZH12

2283

HARUN ÖNLÜ^a, MARIE-CLEMENCE DUCHENE^b, PASCAL HOLLS^b, OZLEM

2284

OSMANAGAOLU^c

2285

2286

^aMuş Alparslan university

2287

^bFaculty of sciences, universite catholique de louvain, institute of life sciences, bbgm lab.

2288

^cAnkara university, faculty of sciences, department of biology, biotechnology units

2289

^e

2290

harunonlu@gmail.com

2291

Lactic acid is a good alternative to petroleum-based plastics because of its many properties. In recent years, production of lactic acid has attracted researchers because of its great potential applications in widespread range of industrial fields. It is the aim of this study to knock-out the gene responsible for L-lactic acid production to obtain strain producing optically pure D-lactic acid to be used in polymer synthesis. **Material-methods:** PCR amplified chromosomal DNA of *Lb. helveticus* OZH12 was used in this study. Disruption and substitution of the *ldhL* gene of YN were carried out using pGIT707 plasmid-based double-crossover homologous integration. The gene expression level of L-lactate dehydrogenase gene was controlled by qRT-PCR. **Result and Conclusions:** Plasmid pGIT707 was transferred to *Lb. helveticus* OZH12 by electroporation. Following transformation the cells were developed in antibiotic containing MRS agar medium and integration was checked by PCR using integration control primers. Positive clones P7, 27b, and 47a were selected and used for the measurement of LDH enzyme activity. Control of gene expression level results demonstrated reduction in *ldh-L* gene expression level and increment in *ldh-D* activity in clone 47a when compared with wild type strain. These results are in parallel with the results obtained from enzyme activity assays and confirm that the lactic acid pathway has changed towards to D-lactic acid production.

2310

2311

Keywords: Lactate Dehydrogenase Gene, Lactic Acid Bacteria, D-Lactic Acid, Knock-Out, Qrt-Pcr

2312

2313

Acknowledgements: Thanks to THE SCIENTIFIC AND TECHNOLOGICAL RESEARCH COUNCIL OF TURKEY (TUBITAK-BİDEB) and Ankara University Scientific Research Project Coordination Unit (AU BAP) for their support.

2314

2315

2316

2317

Poster - 9

2318 **EFFECTS OF ACETYLSALICYLIC ACID ON INFLAMMATION AND**
2319 **OXIDATIVE STRESS IN RATS WITH EXPERIMENTALLY INDUCED**
2320 **TYPE 2 DIABETES MELLITUS**

2321 **SEFA ÇELİK^a, SERKAN ŞEN^b**

2322

2323 ^aDepartment of biochemistry, faculty of medicine, Afyonkocatepe university

2324 ^bAtaturk vocational school of health services, Afyonkocatepe university

2325

2326

sefa_celik@hotmail.com

2327 **Abstract:**

2328 It is believed that adipose tissue-derived molecules contribute to systemic insülin
2329 resistance and to the pathogenesis of most of the metabolic complications of type 2
2330 diabetes mellitus, atherosclerosis, and obesity. Diabetes is considered to be a very
2331 important factor in promoting low density lipoprotein (LDL) oxidation. Stimulation
2332 of lipid peroxidation in LDLs during diabetes is associated with strong oxidation of
2333 cholesterol and glucose during oxidative and carbonyl stress, respectively. These
2334 data describe the relationship between hyperglycemia and atherosclerosis in
2335 diabetic patients. Many studies have shown that inflammation plays a very
2336 important role in the pathogenesis of T2DM. The aim of this study was to determine
2337 the effects of acetylsalicylic acid (ASA) on inflammation and oxidative stress in rats
2338 with experimentally induced T2DM. Obesity was established by body weight gain at
2339 a 5% level and insulin resistance was determined in the rats fed with a high-fat
2340 diet. T2DM was induced by a streptozotocin injection in obese rats. ASA at a dose of
2341 150 mg/kg b.w. was given orally once daily for five weeks. The levels of pro-
2342 inflammatory cytokines, total antioxidant status (TAS), total oxidant status (TOS),
2343 and c-reactive protein (CRP) were analysed. Although TNF- α was increased in the
2344 blood, ASA treatments did not effect the levels of cytokines in the blood and brain
2345 tissue of diabetic rats. Induction of diabetes increased both TOS and TAS in the
2346 blood. ASA treatment also increased blood TOS of diabetics. These results indicated
2347 that diabetes induced inflammation and oxidative stress. It may be said that ASA
2348 treatment at a dose of 150 mg/kg b.w. for five weeks is not sufficient for
2349 amelioration of the inflammation induced by diabetes. **This study confirmed** that
2350 ASA has antioxidant effects. Further studies are needed to investigate the role of
2351 ASA in pro-inflammatory immuno-modulation in diabetes.

2352

2353 **Keywords:** Acetylsalicylic Acid, Inflammation, Oxidative Stress, Type 2 Diabetes

2354

2355

Poster - 10

2356 **THE EFFECT OF CAFFEIC ACID PHENETHYL ESTER (CAPE), A**
2357 **BIOACTIVE COMPOUND OF PROPOLIS EXTRACT, ON ENDOPLASMIC**
2358 **RETICULUM (ER) STRESS-INDUCED APOPTOTIC CELL DEATH IN**
2359 **INS1-B CELLS**

2360

SEFA ÇELİK, AHMET KAHRAMAN, ÖMER HAZMAN, SERKAN ŞEN

2361

2362

Department of biochemistry, faculty of medicine, afyon kocatepe university

2363

2364

sefa_celik@hotmail.com

2365

2366

2367

2368

2369

2370

2371

2372

2373

2374

2375

2376

2377

2378

2379

2380

2381

2382

2383

2384

2385

2386

2387

2388

2389

2390

Caffeic acid phenethyl ester (CAPE), a bioactive component of propolis extract, has many biological and pharmacological properties such as antitumoral, antiinflammatory, and antineoplastic. Endoplasmic reticulum (ER) is the main intracellular organelle responsible for protein folding, translocation and post-translational modification. The ability of the cell to respond to perturbations in ER function or 'ER stress' is critical for cell survival. However, chronic or unresolved ER stress can lead to apoptosis. It is well known that ER stress inducing agents are potential anticancer therapies. In this study, it was aimed to investigate the effect of CAPE on ER stress-induced apoptotic cell death in INS1- β cells in vitro. For this aim, mRNA expression levels of some mediators including CCAAT-enhancer-binding protein homologous protein (CHOP), c-Jun N-terminal kinase (JNK), caspase-12, activating transcription factor 6 (ATF6), B-cell lymphoma (Bcl)-2 and Bcl-2-associated X protein (Bax) involved in ER stress-induced apoptosis were analyzed by Real Time Quantitative PCR in INS1- β cells. After reaching the appropriate number, the cells were incubated with CAPE (10 mmol/L) for 18h. After the incubation, INS1- β cells were harvested for total RNA isolation and quantitative real-time PCR to determine the changes in mRNA expression levels of selected genes following 10 mmol/L CAPE treatment. RNA was extracted after treatment and the fold changes were calculated with β -actin as the internal control. CAPE treatment was able to induce ER stress and apoptosis. Finally, CAPE treatment has resulted in a significant CHOP, JNK, CAS-12 and Bax overexpression, which are also involved in the molecular mechanism of ER stress-induced apoptosis. In addition, antiapoptotic BCL-2 expression was suppressed by CAPE treatment. **In conclusion**, these findings have raised the exciting possibility of targeting UPR components in pancreatic neuroendocrine tumors therapy. It was also concluded that CAPE administration may be exploited as a strategy for insulinoma treatment.

2391

2392

Keywords: INS1- B Cell, Insulinoma, Caffeic Acid Phenethyl Ester, Endoplasmic Reticulum Stress, Apoptosis.

2393

2394

Poster - 11

2395

EFFECT OF HYPEROSMOTIC STRESS ON MINPP1 EXPRESSION

2396

FATİH ERCİ^a, MUSTAFA YÖNTEM^b, BEHİÇ SELMAN ERDOĞDU^c

2397

2398

^aNecmettin Erbakan university

2399

^bdepartment of biotechnology, faculty of science, Necmettin Erbakan university

2400

^cdept. of molecular biology and genetics, faculty of science, Necmettin Erbakan university

2401

2402

ferci@konya.edu.tr

2403

Hyperosmotic stress is a specific state that can lead to cell death by apoptosis. A hyperosmotic environment provides us an applicable perspective to examine the identifications of the novel signaling models. The studies of the MINPP1, which acts as a phosphoinositide 5- phosphatase and phosphoinositide 6-phosphatase and regulates cellular levels of inositol pentakisphosphate (InsP5) and inositol hexakisphosphate (InsP6) and has an important role in bone development, is essential to know what means cells control the signaling activities of `higher' inositol polyphosphates. Here we tried to identify hyperosmotic stress effect of Minpp1 expression in Hela cell line by expecting effects of hyperosmotic stress on mammalian cells can be determined by its intensity and duration. In this study, HELA cells were exposed for different times (15 min and 30 min) and with different amount of sorbitol. After protein extraction, samples were run into sds-page gel. In order to determine the effect of hyperosmotic stress on Hela cell lines, immunoblotting assay was done with western blot technique and related protein bands were detected on X-Ray film. Hela cells showed a significant response change in the expression of the protein in 15 minutes with increased concentrations of sorbitol. Expression of Minpp in Hela was induced by 100mM and 200mM sorbitol while it decreased in 400 mM sorbitol. The range of hyperosmotic stress to which Hela cell was exposed, up to 200 mM, induced expression of Minpp and this range can be thought as sublethal whereas increasing this amount, namely, the concentration of sorbitol is higher than 400 mM can be thought as an amount for starting limitation of expression Minpp. We **conclude** that certain amount of sorbitol and long duration exposure on Hela cell line can lead to apoptosis.

2426

2427

Keywords: Hyperosmotic Stress, Apoptosis, Minpp1

2428

2429

Poster - 12

2430

BIOCHEMICAL AND HAEMOTOLOGICAL MARKERS IN PATIENTS WITH CHRONIC HBV INFECTIONS

2431

2432

BILGEN ÖZDEMİR^a, AYŞE RÜVEYDA UĞUR^b, MEHMET ÖZDEMİR^b

2433

MAHMUT BAYKAN^b

2434

^aKonya numune hospital

2435

^bDepartment of medical microbiology, faculty of meram medicine, Necmettin Erbakan

2436

university, Konya, TURKEY

2437

2438

numunelab@hotmail.com

2439

Chronic hepatitis B virus (HBV) infection is one of the major public health problems all over the world. Pathophysiology of the development of chronic HBV infection remain poorly understood. A better understanding of biochemical markers of chronic HBV infection may be useful in the disease progress. The aim of this study is to investigate the levels of biochemical and hematological markers, and alpha-fetoprotein (AFP), a tumor marker, in patients with chronic hepatitis B infection. **Serum aspartate transaminase** (AST), alanine transaminase (ALT) levels, lymphocyte/leukocyte ratio, and AFP levels of patients with chronic HBV infection were retrospectively investigated in Medical Microbiology Laboratory Meram Medical Faculty Hospital between January and December 2016. HBV DNA levels were determined by using real time PCR assay via an automatic system COBAS AmpliPrep-COBAS TaqMan (Roche, France). AST and ALT levels were measured by the catalytic activity concentration of enzymes according to IFCC (International Federation of Clinical Chemistry). AFP levels were measured by using chemiluminescence assay. Lymphocyte/leukocyte ratios were determined by using Complete Blood Count with 5-part differential assay. **Of 2349** patients with chronic HBV infection, 237 were enrolled to the study based on the cut-off value of 20.000 IU/mL for HBV DNA. AST levels of 56 and ALT levels of 64 patients were higher than normal. Only 30 (12.7%) patients had a high AFP levels except for two pregnant women whose AFP levels were increased due to pregnancy. Fifteen patients had a lower level than normal for lymphocyte/leukocyte ratio, and only one patient had a higher lymphocyte/leukocyte ratio than normal. **A considerable** amount of patients with chronic HBV infection still have a higher level of AST and ALT. Further detailed investigations are needed to clarify the importance of biochemical and hematological markers.

2464

Keywords: Chronic, HBV, AST, ALT, Lymphocyte/Leukocyte Ratio, Afp

2465

Poster - 13

2466

INVESTIGATION OF PARAINFLUENZA VIRUS CAUSED LOWER RESPIRATORY TRACT INFECTIONS IN PEDIATRIC PATIENTS

2467

2468

AYSUN GÖRKEM^a, AYŞE RÜVEYDA UĞUR^a, BAHADIR FEYZİOĞLU^a

2469

MEHMET ÖZDEMİR^a, MAHMUT BAYKAN^a

2470

2471

^aDepartment of medical microbiology, faculty of meram medicine, necmettin erbakan
university, Konya, TURKEY

2472

2473

2474

mbaykantr@hotmail.com

2475

Objective: Parainfluenza viruses are associated with significant morbidity and mortality in infants and children. The aim of this study is to investigate the rate of parainfluenza infections in pediatric patients with lower respiratory tract infections.

2476

2477

2478

Materials and Methods: Nasopharyngeal swab specimens of pediatric patients with the diagnosis of lower respiratory tract infection were collected in Medical Microbiology Laboratory, Meram Medical Faculty Hospital between January 2014 and December 2016. Parainfluenza types 1, 2, 3, and 4 were identified by using two different commercial kits by multiplex real time PCR method. **Results:** One of the four parainfluenza types was identified in 224 of 1983 nasopharyngeal swab specimens of patients aged under 18 years (11.3%). Parainfluenza type 3 was the most common agent among the subtypes (75%) followed by parainfluenza type 4 (15.2%). The rate of parainfluenza type 1 and parainfluenza type 2 were 5.8% and 4%, respectively. Of all pediatric patients, 75.9% were under 5 years of age. When the seasonal distribution was examined, it was found that there was an increase in the rate of parainfluenza virus infections in the end of summer and autumn (65.6% between July and November). **Conclusion:** Parainfluenza type 3 was the most common subtype. Prompt identification of parainfluenza viruses by rapid diagnostic methods would be very helpful for clinicians in the diagnosis and prevention of morbidity and mortality caused by these viruses.

2479

2480

2481

2482

2483

2484

2485

2486

2487

2488

2489

2490

2491

2492

2493

2494

2495

Keywords: Parainfluenza Virus, Pediatric, Multiplex Pcr

2496

Poster - 14

2497

INVESTIGATION OF PARVOVIRUS B19 IGM AND IGG POSITIVITY RATES IN PEDIATRIC HEMATOLOGY PATIENTS

2498

2499

AYSUN GÖRKEM^a, AYŞE RÜVEYDA UĞUR^b, FATMA TAŞBENT^c, MEHMET ÖZDEMİR^d, MAHMUT BAYKAN^e

2500

2501

2502

^aDEPARTMENT OF MEDICAL MICROBIOLOGY, FACULTY OF MERAM MEDICINE, NECMETTIN ERBAKAN UNIVERSITY, KONYA, TURKEY

2503

2504

2505

mbaykantr@hotmail.com

2506

Human parvovirus B19 is a frequent etiologic agent causing erythema infectiosum in children. It has recently been suggested that parvovirus B19 may be latent after infection and cause reactive infections especially in immunosuppressive patients with hematological problems. In this study, we aimed to investigate the parvovirus B19 IgM and IgG positivity rates in sera of patients consulted Pediatric Hematology clinic. **We retrospectively screened** the laboratory results of parvovirus B19 IgM and IgG antibody assays of children < 18 years, who consulted Pediatric in and outpatient clinics at between 2013-2016. Parvovirus B19 IgM and IgG antibodies were investigated in serum samples by using enzyme linked immunosorbent assay method in the Medical Microbiology Laboratory. **Regarding all kinds of pediatric** in and outpatients, the total number of patients from whom parvovirus B19 IgM antibodies were investigated was 2697 and parvovirus B19 IgM antibodies were detected in 168 (6.2%). On the other hand, of all, the number of patients attending Pediatric Hematology clinics was 602 and parvovirus B19 IgM antibodies were detected in 109 (18.1%). A total of 1299 patients were asked for parvovirus B19 IgG antibody and 368 (28.3%) of them were positive for parvovirus B19 IgG. Of all, parvovirus B19 IgG antibody was detected in 244 (25.6%) of the 952 samples collected from Pediatric Hematology clinics. Parvovirus B19 IgM and IgG positivity in samples from Pediatric in and outpatient clinics other than Pediatric Hematology were 2.8% and 35.7%, respectively. Parvovirus IgM positivity in serum samples sent from the Pediatric Hematology in and outpatients was statistically significant compared to those sent from Pediatric clinics other than Pediatric Hematology ($p = 0.0001$). **The higher detection rate of serum parvovirus B19 IgM** positivity in patients under the follow-up of Pediatric Hematology clinics suggests that immune suppression-related virus reactivation or persistence may occur in these patients.

2531

2532

Keywords: Parvovirus B19, Igm, Igg, Child Patient

2533

2534

Poster - 15

2535

INVESTIGATION OF BOKAVIRUS-RELATED LOWER RESPIRATORY TRACT INFECTIONS IN PEDIATRIC PATIENTS

2536

2537

**AYŞE RÜVEYDA UĞUR, AYSUN GÖRKEM, FATMA TAŞBENT,
BAHADIR FEYZİOĞLU, MEHMET ÖZDEMİR**

2538

2539

2540

Department of medical microbiology, faculty of meram medicine, necmettin erbakan
university, konya, turkey

2541

2542

2543

mehmetozdem@yahoo.com

2544

Objective: Human Bokavirus is a newly discovered virus belonging to Parvoviridae family and frequently detected in children's respiratory tract specimens. The aim of the present study is to investigate the frequency of bokavirus in pediatric patients with lower respiratory tract infections. **Materials and Methods:** Nasopharyngeal swab specimens of patients with the diagnosis of lower respiratory tract infection, who are aged < 18 years and followed in various pediatric clinics, were collected between August 2014 and July 2017 in Medical Microbiology Laboratory, Meram Medical Faculty Hospital. Bokavirus was consecutively identified by using two different commercial kits by multiplex real time PCR method. **Results:** As a result of the study, bokavirus was detected in 85 of 1524 pediatric patients (5.6%). Of these 85 patients, 65 were found to be ≤ 5 years old (76.4%). When the seasonal distribution was examined, it was found that the frequency of bokavirus infection was higher in June and July (41.1%). Cough and fever were the most frequent symptoms of the patients (70.5% and 61.1%, respectively). **Conclusion:** The findings of the present study suggest that bokavirus may be a rare but important viral agent causing lower respiratory tract infections especially in children younger than 5 years. On the other hand, since it is a newly discovered virus, further studies are needed on prevalence and risk factors of bokavirus infections.

2562

2563

Keywords: Bokavirus, Pediatric, Multiplex PCR, Prevalance

2564

Poster - 16

2565 **An investigation of blood selenium, copper and zinc levels in sheep**
2566 **at region of Siverek**

2567

MEHMET BATMAZ^a, ILKER CAMKERTEN^b

2568

2569

^aVeterinarian in Şanlıurfa & Health Science Institute, Harran University

2570

^b Dept. of internal medicine, faculty of vet. med. Aksaray University, **Supervisor,**

2571

2572

ilkercamkerten@gmail.com

2573

Background: Due to its critical role in organism, trace elements have a very important place in the life of the creatures. Trace mineral deficiencies in animals are one of the major economic problems in our country as well as in the whole world. Trace element deficiencies that it's frequently seeing are as important as macro- and micro-element deficiencies, as those resulting from infectious and parasitic diseases. In our country, trace element deficiencies are common but its incidence and severity can't be determined completely. According to Turkvet data, as of 2011 there are 1,223,823 head sheep in Şanlıurfa. **In this study,** it was aimed to determine the levels of Se, Cu and Zn elements in sheep breeding in Şanlıurfa-Siverek region in the context of preventive medicine. **Materials and Methods:** In this study, Siverek district was divided into four regions as north, south, east and west in the spring lambing period, and blood samples were collected from 100 sheep randomly selected from a total of 25 sheep from each region. Whole blood samples were transplanted to Elazığ Veterinary Control Research Institute, Se, Cu and Zn analyzes were carried out by Perkin Elmer Aanalyst 800 (HGA Graphite Equipments, USA) with AOAC 2000 and 999.10 method. The SPSS statistical package program was used to evaluate individual data, variance analysis was performed, Duncan multiple comparison test was used to determine the difference between the groups. **DISCUSSION:** In this study, values were determined using Full Blood. Considering that reference values are obtained from serum / plasma for Cu and Zn, in particular, it was also considered that the use of whole blood may be an effect of high blood Cu and Zn levels in our study. **In conclusion,** we consider that the data obtained in this respect, which is very important for humanity, animal husbandry, region and country economy, contributes to the geographical differences and the selenium, copper and zinc levels in the region.

2598

2599

Keywords: blood, selenium, copper, zinc, sheep

2600

Acknowledgements: *This study is the MSc thesis of Mehmet Batmaz, DVM, MSc.*

2601

Poster - 17

2602 **THE POST-HOC POWER ANALYSIS OF FOREST PRODUCTIVITY**
2603 **ATTRIBUTES IN EXPERIMENTAL STUDY IN CENTRAL BOSNIA**

2604 **AZRA ČABARAVDIĆ, AIDA IBRAHIMSPAHIĆ, MIRSADA STARČEVIĆ**

2605

Faculty forestry university Sarajevo

2606

2607

2608

azrac2012@gmail.com

2609 **Forest** productive attributes changes over time in native forests has been
2610 recognized as crucial challenge for management of mixed uneven aged forests in
2611 Bosnia and Herzegovina since middle of the last century. Experimental study has
2612 been carried out on set of experimental plots established in mixed stands on
2613 mountain Igman in central Bosnia. The most important forest productivity
2614 attributes changes based on repeated measures have been monitored over time. The
2615 aim of this research was to conduct the post-hoc power analysis for monitored
2616 forest attributes: basal area per ha (BA), growing stock per ha (GS) and current
2617 annual increment per ha (CAIv). **Here are used repeated measures** conducted on
2618 the 10 experimental plots in two type of mixed stands: fir-spruce and fir-spruce-
2619 beech plots (five plots per each type) measured in five (BA and GS) and four (CIVv)
2620 occasions in periods between 10-20 years. Analyses of variance (ANOVA) within and
2621 within-between repeated measures were applied and power analysis was performed.
2622 ANOVA within forest type over time showed highly significant differences for all
2623 attributes ($\alpha=0.05$, $p<0.001$). Here, power analysis for comparison of stand
2624 attributes resulted in observed high power values ranged from 82% to 99% (very low
2625 risk of Type II errors). Then, ANOVA between two forest types over time showed
2626 different significances for forest attributes ($\alpha=0.05$, $p_{BA}=0.25$, $p_{GS}=0.23$ and
2627 $p_{CIVv}=0.02$). The risks of Type II errors were high for BA and GS (from 66% to 72%)
2628 while conclusions for CAIv could be accepted with very low risk (4%). So, the post-
2629 hoc power analysis of comparisons of stand attributes between forests types found
2630 low power for BA (28%) and GS (34%) and high power for CAIv (96%). **These**
2631 **findings** confirm importance of proper forest species composition planning in mixed
2632 stands related to highest wood productivity and other forest characteristics as
2633 biodiversity.

2634

2635 **Keywords:** Forest Productivity, Mixed Uneven Aged Forest Stands, Experimental
2636 Study, Repeated Measures, Power Analysis

2637

2638

Poster - 18

2639

RIGOROUS IDENTIFICATION OF TISSUE SPECIFIC GENES IN SILICO: REVEALING THE INTERPLAY BETWEEN CANCER SPECIFIC EXPRESSION AND TISSUE SPECIFIC EXPRESSION

2640

2641

2642

HATİCE BÜŞRA KONUK^a, ALPER YILMAZ^b

2643

2644

^aGebze technical university, department of bioengineering, Kocaeli, TURKEY

2645

^byildiz technical university, department of bioengineering, İstanbul, TURKEY

2646

2647

hbkonuk@gtu.edu.tr

2648

Tissue-specific genes are a class of genes whose function and expression are preferred in one or several tissues, specifically. Identification of tissue-specific genes is essential for discovery of tissue-specific targets and understand molecular mechanism. In this study, we aimed to identify tissue-specific genes in more rigorous fashion, examine intersection between differentially expressed genes (DEGs) in various cancers and tissue specific genes, interpret tumor heterogeneity, reveal tissue specific molecular targets for different cancers and elucidate role of tissue specific genes in biological processes. **Gene expression data**, derived from five large RNA-Seq projects, spanning 96 different human tissues was retrieved from ArrayExpress and ExpressionAtlas. We developed an algorithm using R programming language and rigorous identification of tissue specificity was calculated by integrating tau score and statistical distance methods. DEGs of 18 different cancers were taken from BioExpress and we investigated intersection of cancer DEGs with corresponding tissue-specific genes. Tissue-specific genes and overlapping genes were analyzed with DAVID for GO terms, related diseases and pathways associated with each gene. In addition, tissue-specific unusual transcripts from RNA-Seq data is under investigation by our group. **We successfully** assigned genes to multiple tissues for specificity using novel approach. Functional metadata of these genes showed that they are associated with diseases related with corresponding tissue. After intersection, overlapping genes was found to be quite small, it can be significant for targeted treatment of solid tumors. If we investigate subtypes of one cancer, overlapping genes can be different. These result shows us mechanism of subtypes are different. More importantly, we observed that many DEGs in one cancer might contain genes specific to other tissues which might be due to tumor heterogeneity. These results suggest that cancer tissues might be heterogeneous and cancer networks might have complex interplay with tissue-specific genes.

2675

2676

Keywords: Tissue-Specific Genes, RNA-Seq, Tau Score, Statistical Distance, Tumor Heterogeneity

2677

2678

Poster - 19

2679

GENERATING LOSSLESS COMPRESSION OF GENOME SCALE K-MER FREQUENCY TABLE AS RASTER IMAGE

2680

2681

2682

HATICE BÜŞRA KONUK^a, ALPER YILMAZ^b

2683

2684

^aGebze technical university, department of bioengineering, Kocaeli, TURKEY

2685

^bYıldız technical university, department of bioengineering, İstanbul, TURKEY

2686

2687

hbkonuk@gtu.edu.tr

2688

K-mer frequency of any DNA sequence is calculated by counting occurrences of all possible substrings of length k . The k -mer frequency of genome or next generation sequencing data is an invaluable tool to gain insights about the DNA sequence and its grammar. For genomes, k -mer counts can be used for motif discovery, classification and alignment-free comparison of multiple genomes. For short reads, k -mer counts are used for quality check, diagnosis, error correction and assembly. The initial step k -mer counting requires storage of frequency tables which tend to get bigger by increasing length of k . In this study we propose a method for lossless compression of k -mer data which is expected to simplify and facilitate storage and analysis of k -mer data. **In a raster image**, such as PNG, each pixel has two components; coordinate and color. We implemented Chaos Game Representation (CGR) to map k -mers to coordinates and k -mer occurrence was mapped to RGB color via bit-level operations. CGR maps can be divided and labeled according to the corresponding substring, each substring is mapped onto a sub-square, creating a fractal-like structure. Basically, the whole set of frequencies of the k -mers found in each genomic sequence are displayed in the form of a single image in which each pixel is associated with a specific k -mer and its occurrence. **As result**, file size has been reduced by approximately 10 times compared to plain text and reduced 5 times compared to binary storage (Jellyfish). Storage of k -mer data as image will not only save storage space but also facilitate genomic analysis in a manner previously not implemented. Image related algorithms can be used to process, analyze and manipulate collection of images representing genomic or next-generation sequencing data k -mer signatures.

2711

2712

Keywords: Genome, K-Mer, Lossless Compression, Chaos Game Representation, Png

2713

2714

Poster - 20

2715

DESIGN AND PRODUCTION OF A LAB-SCALE SOLAR BIOREACTOR

2716

VILDAN ERCI^a, DIDEM ÖZÇİMEN^b, İBRAHİM İŞILDAK^b

2717

2718

^aselcuk university

2719

^byildiz technical university

2720

2721

vildanerci@selcuk.edu.tr

2722

Microalgae have a great biotechnological potential in the production of valuable raw materials used in silage, food, agriculture, cosmetics and pharmaceutical industries as well as biotechnological productions. For this reason, it is an important issue to base microalgae production on a biotechnological basis. In both open and closed microalgal culture systems, light source and light intensity are critical factors affecting the phototrophic growth performance of microalgae. **In this study**, a solar bioreactor design was carried out in order to provide controllable, high efficiency and low cost production. The designed bioreactor was built on a laboratory scale solar system for microalgae growth. The solar system consists of 3 cylindrical tubes, windscreen wipers on the tubes, fresnel lenses and fiber optic cables.. Fresnel lenses enrich the sun light and transferred into the bioreactor by fiber optic cables. In the system, each fresnel lens is connected to a fiber optic cable so that the sunlight is delivered homogeneously through the fiber optic cables to the area where microalgae are present and the daylight required for microalgae for photosynthesis can be provided. Fiber optic cables transfer the light to the cylindrical tubes where they are placed, and illuminate inside of the reactor homogeneously in which needed by the microalgae. **We can conclude that** the microalgae, which are thought to be the nutrients of the future, are produced in a controlled and rapid manner, making this system valuable. In addition, it is revealed that the problems such as the high costs of the production and contamination can be eliminated by this system. Finally, it would be possible to get about 4-5 times higher quality product with 100 lt closed bioreactor used in this study than an open system.

2745

2746

Keywords: Microalgae, Solar Bioreactor, Fiber Optic Cable.

2747

2748

Poster - 21

2749 **Evaluation of ischemia modified albumin levels in subclinical**
2750 **mastitis cows before and after the vaccination**

2751 GAYE BULut^a, GÜZİN CAMKErten^b, ÖZCAN EreI^c

2752

2753 ^a Dept. of Obstetrics and Gynecology, Faculty of Veterinary Medicine, ^b Dept. Veterinary,
2754 Vocational School of Technical Sciences, Aksaray University Aksaray

2755 ^c Ankara Atatürk Training and Research Hospital Biochemistry Laboratory, Ankara
2756 TURKEY
2757

2758

oguzalperen@hotmail.com

2759 **Introduction:** Mastitis is the inflammatory response of the mammary gland tissue
2760 to physiological and metabolic changes, traumas, and allergies and, most
2761 frequently, to injuries caused by various microorganisms. The inflammation formed
2762 during these events leads to a decrease in milk production and to some changes in
2763 its structure. Also, it is seen that blood values changed in all types of inflammatory
2764 response of the mammary gland. **This investigation** was implemented to assess
2765 the IMA that is a marker for inflammatory processes in dairy cattle with subclinical
2766 mastitis by vaccination. **Methodology:** A fifty-one dairy cow were included in the
2767 study. SCC: Tank milk sample was collected only one and transferred into 200 ml
2768 plastic tubes, and SCC were measured from fresh milk samples by Bentley -Merkim
2769 B-150 M150 instruments. A total of 51 blood sample before and after mastitis
2770 vaccination. 37 of the sera samples were taken and were done biochemical
2771 evaluation for IMA. The others were hemolysis. Groups were compared using
2772 Student's t-test. **Results and Discussion:** In our study, there was no statistically
2773 significant change in classic IMA. But There was statistically significant change in
2774 Modified IMA ($p < 0.05$). Elevated ischemia modified albumin (IMA) levels have been
2775 reported as a result of ischemia, which is due to increased inflammatory processes.
2776 **We determined** increase of modified IMA as markers of inflammation. Ischemia-
2777 induced oxidative stress may start proinflammatory processes. On the other hand,
2778 IMA may come forward as a marker for inflammatory processes. **Summary:** Our
2779 study will be the first report in the literature. The knowledge oxidative defense
2780 mechanisms of animals will serve as a guiding principle for future studies.

2781

2782 **Keywords:** Sub clinical mastitis, ischemia modified albumin, vaccination, cow

2783

2784

Poster - 22

2785 **Diversity in mixed uneven aged forest of beech, fir and spruce on**
2786 **mountain Igman**

2787 **AIDA IBRAHIMSPAHIĆ, AZRA CABARAVDIC, MIRSADA STARCEVIC**

2788

2789 Faculty of forestry university Sarajevo

2790

2791 azrac2012@gmail.com

2792 **With the aim of understanding** of complicated laws regarding the process of trees
2793 and stands' growth, and precise prediction of their development within up-to-date
2794 researches, most of attention is focused on spacial configuration and individual
2795 differences. During structural analysis numerous indicators are determined which
2796 are, apart from horizontal and vertical description of stems' structure also useful for
2797 diversity descriptions of species, positions and dimensions on stand level or nearby
2798 to referential trees. **Based on data** collected on eight permanent experimental plots
2799 in uneven aged mixed forests of beech, fir and spruce on mountain Igman the
2800 determined sizes of Shannon-Weaver (H' : 0,68-1,12) and Simpson (λ : 0,34-0,57)
2801 indices indicate high level of species diversity. The determined sizes of Clark and
2802 Evans index (CE: 1,10-1,40) indicate that the horizontal spatial distribution of trees
2803 strives for being regular. Vertical one, according to the sizes of species profile index
2804 (A: 1,24-1,95) and index by Kotar (B: 0,16-0,40), are even for five experimental plots
2805 and uneven for other three. Mixing of tree species on experimental plots, according
2806 to the M_i index sizes (fir 0,24-0,61; spruce 0,28-0,75; beech 0,18-0,76;), are
2807 different. Mixing by individual trees, tree species of a smaller portion are individually
2808 mixed with tree species of a bigger portion, trees of present species are in groups
2809 and trees of present species individually and in groups. **Detemined sizes** of index
2810 of size differentiation (T: 0,31-0,44) indicate that diameters of referential and
2811 neighbouring trees are in such relation that the smaller diameter makes about 60-
2812 70% of the bigger one. According to sizes of index of dominance (U: fir 0,31-0,67,
2813 spruce 0,38-0,70, beech 0,33-0,73) referential trees of fir are dimensionally
2814 relatively more dominant than ones of another tree species on four experimental
2815 plots, referential trees of spruce are more dominant on two plots as well as beech.

2816

2817 **Keywords:** mixed uneven aged forest, species diversity, positions diversity,
2818 dimensions diversity

2819

Poster - 23

2820

PAIR HOUSING FOR CALVES

2821

MUSTAFA ÇAM^a, ŞEREF İNAL^a

2822

2823

^aSelçuk university veterinary faculty animal science department

2824

2825

mustafa0609@hotmail.com

2826

Calves usually rear at individual hutches till weaning in dairy farms. But it should be known that socialization has some beneficial effects on calves with regard to stress, behaviour, performance, appetite. Also it's known that rearing calves with small groups doesn't have serious adverse effect on their health and comfort. Recently it has been done some researches about calf housing systems affected calf socialization, comfort and health. So in **this presentation** it was aimed to give some information about rearing calves with pair housing.

2827

2828

2829

2830

2831

2832

2833

2834

Keywords: Pair, Calf, Housing, Dairy

2835

2836
2837
2838
2839
2840
2841
2842
2843
2844
2845
2846
2847
2848
2849
2850
2851
2852
2853
2854
2855
2856
2857
2858
2859
2860
2861
2862

interim

FULL TEXT PRESENTATIONS

2863

Oral - 56

2864 **The Effect of Spent Mushroom Compost on the Growth, Mineral**
2865 **Nutrition and HeavyMetal Status of Lettuce (*Lactuca sativa* L.) and**
2866 **Spinach (*Spinaceae oleraceae* L.)**

2867

BÜLENT TOPCUOĞLU, M. KUBILAY ÖNAL

2868

2869 Akdeniz University Vocational School of Technical sciences, 07058 Antalya, TURKEY

2870

*Corresponding author: btoglu@akdeniz.edu.tr

2871

2872 **Abstract**

2873 This research was carried out to determine the effects of spent mushroom compost (SMC) as
2874 an organic material source on the growth, plant nutrient and heavy metal content of lettuce
2875 and spinach plants grown in greenhouse soil. Lettuce and spinach plants were grown in pots
2876 containing different amounts of SMC (corresponding to 0, 20, 40, 60 and 80 T/ha, as dry
2877 weight basis). The effects of SMC on plant dry matter and N, P, K, Ca, Mg, Fe, Zn, Cu, Ni,
2878 Cd, Pb and Cr contents of lettuce and spinach plants were determined.

2879 SMC application increased soil organic matter and EC values but decreased pH value. No
2880 changes in soil mineral and heavy metal content were detected. SMC applications caused
2881 statistically important effects on dry matter yield, and N, P, K, Fe and Zn contents both in the
2882 lettuce and spinach plants. SMC applications increased yield until to 60 ton/ha of SMC
2883 application, but highest application rate of SMC compost depressed plant growth only in
2884 lettuce. Spinach plant was more tolerated to higher SMC levels than lettuce. All spent
2885 mushroom compost treatments resulted high mineral content both in lettuce and spinach.

2886 No important changes in heavy metals except Fe and Zn were detected in all plants by SMC
2887 applications. All metal concentrations were ranged in background levels, below the
2888 phytotoxic maximum limits and no metals exceeded referenced food codex values by the
2889 SMC applications. This research showed that SMC could be applicable in vegetable
2890 cultivation at the agronomic rates without heavy metal contamination concern but salinity
2891 defects in salt sensitive plants should be taken into consideration.

2892

2893 **Key Words:** Spent Mushroom Compost, Mineral Nutrition, Heavy Metals, Lettuce, Spinach

2894

2895 **Mantar compost atığının marul (*Lactuca sativa* L.) ve ıspanak**
2896 **(*Spinaceae oleraceae* L.) bitkilerinde gelişme, mineral beslenme ve**
2897 **ağır metal içerikleri üzerine etkisi**

2898 **Özet**

2899 Bu araştırma bir organik madde kaynağı olarak mantar compost atığı'nın (MKA) serada
2900 yetiştirilen marul ve ıspanak bitkilerinde gelişme, bitki besin ve ağır metal içerikleri üzerine
2901 olan etkilerini belirlemek amacıyla yürütülmüştür. Marul ve ıspanak bitkileri değişik
2902 miktarlarda MKA içeren (kuru madde bazında 0, 20, 40, 60 ve 80 T/ha) saksılarda
2903 yetiştirilmiştir. MKA'nın marul ve ıspanakta kuru madde ve N, P, K, Ca, Mg, Fe, Zn, Cu, Ni,
2904 Cd, Pb ve Cr içerikleri üzerine etkileri belirlenmiştir.

2905 MKA uygulaması toprak organik madde ve EC değerlerini arttırmış ancak pH değerini
2906 düşürmüştür. Toprak mineral ve ağır metal içeriğinde önemli değişiklik saptanmamıştır.
2907 MKA uygulamaları her iki bitkide kuru madde ile N, P, K, Fe ve Zn içeriklerini arttırmıştır.
2908 MKA uygulamaları ürün miktarını 60 ton/ha uygulama düzeyine kadar arttırmış, fakat
2909 MKA'nın en yüksek uygulama düzeyinde marul bitkisinde gelişim olumsuz etkilenmiştir.
2910 Ispanak bitkisi yüksek MKA uygulamalarında marul bitkisine göre daha fazla tolerans
2911 göstermiştir. Bütün MKA uygulamaları her iki bitkide daha yüksek mineral içeriği
2912 sağlamıştır.

2913 MKA uygulamalarıyla her iki bitkide Fe ve Zn dışında ağır metal içeriklerinde önemli
2914 değişiklikler kaydedilmemiştir. MKA uygulamalarıyla bütün metal konsantrasyonları bazal
2915 düzeylerde ve de fitotoksik maksimum sınır değerlerin altında saptanmış ve hiç bir metal
2916 önerilen gıda kodeks sınır değerlerini aşmamıştır. Bu araştırma MKA'nın agronomik
2917 oranlarda sebze üretiminde ağır metal kontaminasyon endişesi olmaksızın kullanılabileceğini
2918 ancak tuza hassas bitkilerde kullanımında tuzluluk zararlarının dikkate alınması gerektiğini
2919 göstermiştir.

2920

2921 **Anahtar kelimeler:** Mantar compost atığı, mineral beslenme, ağır metaller, ıspanak, marul

2922

2923 **Introduction**

2924 The use of Spent mushroom compost (SMC) is become to widespread in horticultural
2925 practices as a cheap and abundant waste organic material in mushroom cultivation industry
2926 areas alternative to manure. As in manure applications, SMC applications to soils result a
2927 number of beneficial effects. SMC contains valuable plant nutrients and organic matter that

2928 can improve soil fertility. It can supply plant nutrients to the crop and thus replace inorganic
2929 fertilizer. Trials have shown that it is an excellent source of phosphorus, potassium and trace
2930 elements but needs supplementation with nitrogen for best results. Plant nutrient value of the
2931 SMC is examined by most of the researchers and it is indicated that the nutrient combination
2932 is similar to soil arrangements, based on the organic wastes, applied to the agricultural areas
2933 routinely such as cattle manure and compost (Kütük et al., 1998).

2934 The phytonutritive capacity of compost has often been demonstrated to be analogous to that
2935 of manure; the same level of productivity, both quantitatively and qualitatively, can be
2936 maintained by replacing manure with compost (Beyca et.al.,1993). However, SMC often
2937 contains high salt levels remaining from the fertilizer materials applied during mushroom
2938 cultivation. High or excessive soluble salts may be injurious to plants (Anonymous, 1995).

2939 In this research, the effects of SMC applications to soil on soil chemical characteristics and
2940 soil metals and on plant nutrient contents and heavy metal accumulations in the lettuce and
2941 spinach plants those have different toleration to salinity were examined.

2942

2943 **Materials and methods**

2944 A pot experiment was carried out in the greenhouse and lettuce and spinach plants those
2945 have different toleration to salinity (FAO, 1976) were grown in soil treated with SMC. SMC
2946 material was collected from mushroom cultivation plant in Korkuteli representative of the
2947 major mushroom growing area of Turkey. After collecting the fresh SMC, it was awaited in
2948 open air for 6 months, air dried, and applied to the soil.

2949 In the experiment, plastic pots containing 10 kg soil were used. After 6 months open air
2950 treatment, SMC was applied to experimental soil as an oven-dry basis, corresponding at the
2951 following rates:

2952 - SMC₀: no SMC application (control treatment)

2953 - SMC₂₀: 20 ton/ha of SMC

2954 - SMC₄₀: 40 ton/ha of SMC

2955 - SMC₆₀: 60 ton/ha of SMC

2956 - SMC₈₀: 80 ton/ha of SMC

2957

2958 Pots were arranged in a completely randomized design with four replicates in the greenhouse.
2959 Before beginning of experiment, all treatments received supplemental fertilization at a rate of
2960 150, 50 and 100 mg kg⁻¹ of N, P and K, respectively.

2961

2962 Seedlings of lettuce were grown on peat-perlite mixture and were transplanted as two plants
2963 per pot. Spinach seeds were sowed to pot soil and after the germination, two spinach
2964 seedlings were stayed on the pot. All pots were located in the greenhouse under controlled
2965 climatic conditions. Pots were maintained around field capacity water tension by daily
2966 watering with distilled water. Sixty days after transplanting of lettuce and hand thinning of
2967 spinach, plants were harvested by cutting on the soil surface. Plant samples were washed by
2968 distilled water, dried at 65 °C for 48 h and then ground. In dried leaf samples total N were
2969 determined by Kjeldahl method. Plant tissues were digested in aqua regia (1:3 HNO₃/HCl). In
2970 wet ashed leaf samples total P were determined by molybdophosphoric yellow colour method,
2971 total K, Ca, Mg, Fe, Zn, Mn, Cu, Cd, Ni and Pb were determined by ICP-MS under optimized
2972 measurement conditions.

2973
2974 ANOVA procedures for a randomized complete block design and least significant differences
2975 (LSD) at P<0.05 of data were analyzed by SPSS-16 statistical program.

2976

2977 **Results and Discussion**

2978 *Soil and SMC chemical properties and heavy metal contents*

2979 Greenhouse soil and SMC general analytical characteristics are shown Table 1. The heavy
2980 metal contents of untreated greenhouse soil and SMC are well within the accepted normal
2981 range of values. A comparison of metal contents of SMC with that of untreated soil showed
2982 that the metal concentration Fe, Mn, Ni, Pb and Cr of SMC were not present in greater
2983 concentrations than in the soil. Zinc, Cu and Cd concentration of SMC was found greater. The
2984 heavy metal concentration of SMC is below the referenced levels indicated by the EU (CEC,
2985 1986) for the agricultural use of waste organic material (sewage sludge).

2986 At the end of experiment, changes in soil chemical parameters and heavy metal contents after
2987 treatment of SMC were presented in Table 2. Soil pH, EC and organic matter values were
2988 importantly changed by SMC applications. SMC applications increased EC value and organic
2989 matter content but decreased pH value of soil. All heavy metals were found in referenced
2990 heavy metal limits and no changes were detected in heavy metal contents in the soil by SMC
2991 applications. Increased EC values by SMC applications could be caused by high salt
2992 concentration of SMC. This shows us that salt content is an important factor for a sustainable
2993 SMC application and should be considered in horticultural cultivation before treatment.

2994

2995

2996 Table 1. The analytical characteristic of the experimental soil and SMC, and their pollutant limits.

Parameters	Soil	Limit values in soil ¹ (pH<6-pH>6)	SMC	Limit values in organic materials ¹
Texture	Loam		-	
pH- H ₂ O (1:5 w/v)	7,62		6.74	
EC. (1:5 v/w) dS m ⁻¹	0.205		7.85	
CaCO ₃ , %	12,25		-	
Total N, %	0.21		2.35	
Organic Matter, %	2,76		68,21	
Fe, mg kg ⁻¹	12200	nls	387	nls
Mn, mg kg ⁻¹	460	nls	224	nls
Zn, mg kg ⁻¹	123	150-300	672	2500-4000
Cu, mg kg ⁻¹	32	50-140	55	1000-1750
Ni, mg kg ⁻¹	39	30-75	27	300-400
Pb, mg kg ⁻¹	27	50-300	6	750-1200
Cd, mg kg ⁻¹	0,55	1-3	0,62	20-40
Cr (total), mg kg ⁻¹	41	60-100	12	1000-1750

2997 *: Below detection limit; (< 0.02 mg kg⁻¹); ¹: Total concentrations (mg kg⁻¹ dry wt), (C.E.C., 1986); nls: no limitation set

2998

2999 *Plant yield and heavy metal contents*

3000 The dry matter yield and the concentrations of N, P, K, Ca, Mg, Fe, Mn, Zn, Cu, Ni, Pb, Cd
3001 and Cr in the leaves of lettuce and spinach plants treated with SMC are presented in Table 3
3002 and Table 4, respectively. The effect of spent mushroom compost on the dry matter yield of
3003 lettuce and spinach were found statistically important and SMC increased dry matter yield in
3004 both of plant. The best result for dry matter yield in both plants as regard to productivity was
3005 obtained at 60 ton/ha SMC applications. Dry matter yield decreased in lettuce plant at 80
3006 ton/ha of SMC applications, possibly by high salinity. Whereas at 80 ton/ha of SMC
3007 applications, no changes in dry matter in spinach plant was observed. This may be possibly
3008 due to high toleration of spinach in saline conditions (Table 1 and Table 2).

3009 Plant N, P, K, Fe and Zn contents in both plant were found statistically important. N, P, K, Fe
3010 and Zn contents in both plant were increased by SMC applications, however no important
3011 changes in other nutrients and metal elements were detected (Table 3 and Table 4). Chong et
3012 al. (1991), Wisniewska and Pankiewicz (1989) reported similar findings in their SMC studies.
3013 Mineral and heavy metals except P and K in Spinach plant were detected higher than lettuce
3014 plant. As can be seen in Table 2, nitrogen content of lettuce and spinach was below the
3015 background level. Nitrogen content was increased by SMC applications. This increase was
3016 attributed to increased N availability of SMC (Rhoads and Olsen, 1995; Kutuk et al., 1999).

3017 Although no changes were observed in soil Fe and Zn concentration by SMC applications,

3018 plant Fe and Zn contents increased by SMC treatments. This may be possibly caused by
3019 higher Zn content of SMC and the decreasing pH value caused by SMC application to soil.
3020 All metal concentrations were mostly ranged in background levels and below the phytotoxic
3021 maximum limits. SMC applications to soil did not resulted high metal accumulation in both
3022 plant and metals in both plant were not exceeded the referenced metal limits.

3023

3024 **Conclusion**

3025 The findings support that the SMC has positive effects on the growth of lettuce and spinach
3026 plants. It can be concluded from this perspective that SMC material can be usable without
3027 heavy metal contamination concern in horticultural practices. For this reason, the use of waste
3028 SMC may be an alternative safe organic material source in vegetable production, especially in
3029 organic farming practices. However, in the long-term SMC applications, high EC values of
3030 SMC and its risks on the soil salinity and plant growth should be considered.

3031

3032 **References**

3033 **Anonymous, (1995).** Guidelines for Compost Quality. The Canadian Council of Ministers of
3034 the Environment.

3035 **Beyca, J., De Chant, L., Conditt, M, Jones, B., (1993).** Wet bag composting trial yields
3036 promising results. *Biocycle*, 34(4): 72-75.

3037 **C.E.C. (Council of the European Communities) (1986).** Directive of 12 June 1986 on the
3038 protection of the environment, and in particular of the soil, when sewage sludge is used in
3039 agriculture (86/278/CEE). *Official Journal of the European Communities*, L181, 6-12.

3040 **Chong,C., Cme, R.A., Rnker, D.L., Allan, O.B. (1991).** Growth and mineral nutrient of
3041 containerized woody species in media amended with spent mushroom compost. *J. of the*
3042 *Amer. Soc. for Hort. Sci.*, 116(2):242-247.

3043 **FAO, (1976).** Water Quality for Agriculture. Irrigation and Drainage Paper, No: 29, Rome.

3044 **FAO/WHO (2007).** Codex alimentarius commission, Joint FAO/WHO Food Standards
3045 Programme, Alinorm 7/30/13, thirtieiht session, Roma, Italy, 2-7 July, 2007.

3046 **Jones, J.R, Wolf, B, Mills, H.A. (1991).** Plant Analsis Handbook. Micro Macro Publishing,
3047 Inc.

3048 **Kabata-Pendias, A., Pendias, H. (2000).** Trace elements in soils and plants. CRC Press,
3049 Boca Raton, FL.

3050 **Kütük, C., Topcuoglu, B., Caycı, G. (1998).** The effect of different growing media on
3051 growth of croton (*codiaeam variegatum* ‘Petra’) plant. *M. Şefik Yeşilsoy International*

- 3052 Symposium on arid region soil, September 21-24, 1998, Menemen İzmir, Turkey
- 3053 **Kütük, C., Topcuoğlu, B., Demir, K. (1999).** Effects of different organic matters applied to
3054 soil on yield and some quality factors and mineral contents in spinach plants. Akdeniz
3055 University Journal of the Faculty of Agriculture, 12, 31-36.
- 3056 **Rhoads, F.M., Olsan, S.M. (1995).** Crop production with mushroom compost. Proceedings
3057 of Soil and Crop Sci. Soc. of Florida, 54:53-57.
- 3058 **Wisniewska, G.H., Pankiewicz, T. (1989).** Evaluation of the suitability of spent mushroom
3059 substrate for tulip cultivation. Prace Instytutu Sadownictwa kwiaciarnictwa wskerniewicack,
3060 14:7-13.

interim

361 Table 2. Changes in soil chemical parameters and heavy metal contents after treatment with SMC.

Treatments	pH (1:5 w/v)	EC (1:5 v/w) dS m ⁻¹	Organic matter %	Fe mg kg ⁻¹	Mn mg kg ⁻¹	Zn mg kg ⁻¹	Cu mg kg ⁻¹	Ni mg kg ⁻¹	Pb mg kg ⁻¹	Cd mg kg ⁻¹	Cr mg kg ⁻¹
SMC ₀	7,62	0,37	2,76	12187	455	672	32	39	25	0,55	41
SMC ₂₀	7,62	0,55	2,76	12186	462	680	32	35	28	0,55	39
SMC ₄₀	7,59	2,33	2,82	12200	459	678	32	39	27	0,55	41
SMC ₆₀	7,59	3,85	2,83	12179	445	688	34	42	28	0,55	44
SMC ₈₀	7,60	4,74	2,85	12200	462	678	32	39	27	0,55	41
Significancy	*	**	*	ns	ns	ns	ns	ns	ns	ns	ns

362 **: P<0.01, *: P<0.05, ns: no significance, ¹: C.E.C (1986); nls: no limitation set.

363

364 Table 3. Plant yield, nutrients and heavy metal contents in dry matter of spinach plant treated with SMC.

Treatments	Dry matter g/pot	N	P	K	Ca	Mg	Fe	Mn	Zn	Cu	Ni	Pb	Cd	Cr
		%						mg kg ⁻¹						
SMC ₀	33 c	3.24 d ¹	0,277 d	2.14 c	3,14	0,82	62 b	56	33b	7	1,1	3,0	0,25	0,04
SMC ₂₀	44 b	3.67 c	0,315 c	2.31 b	3,21	0,88	66 b	62	34 b	7	0,8	2,7	0,25	0,06
SMC ₄₀	57 a	4.14 b	0,335 b	2.43 b	2,99	0,78	71 b	77	44 b	9	1,2	2,9	0,30	0,07
SMC ₆₀	67 a	4.67 a	0,344 a	2.51 a	3,18	0,85	78 a	79	58 a	8	1,1	3,1	0,25	0,05
SMC ₈₀	61 a	4.66 a	0,355	2.58 a	3,14	0,92	82 a	78	62 a	9	1,2	3,2	0,30	0,07
Significancy	**	**	**	*	ns	ns	*	ns	*	ns	ns	ns	ns	ns
Background levels ²		4-5	0,4-0,6	6-7	2,3-3,5	0,5-08	60-200	30-250	25-250	5-25	-	-	-	-
Phytotoxic levels ³		-	-	-	-	-	-	-	100-400	20-100	10-100	30-300	5-30	5-30
Food codex limits ⁴							nls	nls	<20	<10	<10	<2	<0,05	<1,3

365 **: P<0.01, *: P<0.05, ns: no significance, ¹: Means within an amendment followed by the same letter are not significantly different at the 005 level, ²: Jones et al (1991); ³: Kabata-Pendias (2000)

366 ⁴: Limitations for fresh vegetables FAO/WHO (2007).

367

368

369

370

371

072 Table 4. Plant yield, nutrients and heavy metal contents in dry matter of lettuce plant treated with SMC.

Treatments	Dry matter g/pot	N	P	K	Ca	Mg	Fe	Mn	Zn	Cu	Ni	Pb	Cd	Cr
		%						mg kg ⁻¹						
SMC ₀	55 c	2,88 d ¹	0.425 d	3,22 c	2,74	0,49	79 b	67	24 b	5	0,8	3,1	0,30	0,05
SMC ₂₀	62 b	2,96 c	0.459 c	3,53 b	2,81	0.51	75 b	71	23 b	5	0,8	3,2	0,35	0,07
SMC ₄₀	73 a	3,15 b	0.538 b	3,45 b	2.88	0.59	88 a	79	27 b	5	0,9	2,8	0,35	0,05
SMC ₆₀	68 a	3,25 a	0.564 a	3,88 a	3,15	0.62	90 a	77	32 a	7	0,8	3,3	0,30	0,06
SMC ₈₀	57 b	3,32 a	0.574 a	3,77 a	2,78	0,58	98 a	81	35 a	7	0,9	3,0	0,35	0,07
Significancy	**	**	**	*	ns	ns	*	ns	*	ns	ns	ns	ns	ns
Background levels ²		4-5	0,4-0,6	6-7	2,3-3,5	0,5-08	50-100	15-250	25-250	8-25	-	-	-	-
Phytotoxic levels ³		-	-	-	-	-	nls	nls	100-400	20-100	10-100	30-300	5-30	5-30
Food codex limits ⁴							nls	nls	<20	<10	<10	<2	<0,05	<1,3

073 **; P<0.01, *, P<0.05, ns: no significance, ¹: Means within an amendment followed by the same letter are not significantly different at the 005 level, ²: Jones et al (1991); ³: Kabata-Pendias (2000)

074 ⁴: Limitations for fresh vegetables FAO/WHO (2007).

075

076

Plant nutrient status of sour cherry (*Prunus Cerasus L.*) cultivars grown in Aegean region of Turkey

BÜLENT TOPCUOĞLU, M. KUBILAY ÖNAL

Akdeniz University Vocational School of Technical sciences, 07058 Antalya, TURKEY

*Corresponding author : btoglu@akdeniz.edu.tr

Abstract

Keywords: Cherry cultivars, Mineral status, Genotypic variability

Türkiye'nin Ege bölgesinde yetişen Vişne çeşitlerinin (*Prunus Cerasus L.*) bitki besin durumları

Özet

3109 Bu çalışma Türkiye'nin Ege bölgesinden toplanan 29 vişne çeşidinde bitki besin içeriklerini ve
3110 onların toprak karakteristikleri ile ilişkilerini belirlemek amacıyla yürütülmüştür. Bu çalışmada
3111 kullanılan vişne çeşitlerinin yaprak ve meyvelerinde biriken toplam N, P, K, Ca, Mg, Fe, Zn, Mn ve
3112 Cu içerikleri belirlenmek suretiyle mutlak gerekli bitki besin elementlerin yaklaşık alımları tahmin
3113 edilmiştir. İncelenen bütün bitki besin maddelerinin vişne çeşitleri arasında önemli farklılık
3114 gösterdiği belirlenmiştir. Vişne yapraklarında Zn dışındaki bütün elementlerin konsantrasyonları
3115 önerilen sınır değerleri içinde saptanmıştır. Vişne meyvelerinde bütün mineral besin içerikleri 1507
3116 ve 1525 vişne çeşitlerinde en yüksek düzeyde kaydedilmiştir. Düşük organik madde ve hafif alkalın
3117 reaksiyonlu toprak koşullarında vişne çeşitlerinin çoğunda Zn içerikleri noksanlık düzeyinde ve
3118 birçok vişne çeşidinin yapraklarında ise N içerikleri düşük düzeyde belirlenmiştir. İncelenen vişne
3119 çeşitlerinin kültüre alınmasında N ve Zn elementleri gübreleme programının önemli bir bileşeni
3120 olarak kaydedilmiştir. Vişne çeşitlerinde yaprak mineral besin içeriklerinin meyve dokusundaki
3121 mineral içeriklerinden daha yüksek olduğu saptanmıştır. Vişne genotipi 1512, yaprak ve meyve
3122 dokularında bütün besin içeriklerini diğer genotiplerden daha düşük düzeyde içermiştir. Aynı toprak
3123 koşullarında yetişen vişne çeşitleri arasında 1515 ve 1530 vişne çeşitlerinin yaprak dokularında
3124 mineral içerikleri diğer çeşitlerden daha yüksek düzeyde bulunmuştur. Bu veriler tarımsal
3125 biyoçeşitliliğin korunması ve vişne çeşitlerinin beslenmesi konularında gelecekte yapılacak
3126 çalışmalar için potansiyel bilgiler sunmaktadır.

3127
3128 **Anahtar kelimeler:** Vişne çeşitleri, mineral içerikler, genotipik çeşitlilik

3130 **Introduction**

3131 Sour cherry is widely planted in the Central Anatolia and Aegean Region of Turkey. These districts
3132 have the best climatic factors, such as brilliant illumination, low temperature and great difference in
3133 temperature between day and night. A few studies (Vesterheim 1973; Christersen 1978; Bekefi *et*
3134 *al.* 2000; Cordeiro-Rodrigues *et al.* 2000) have revealed wide variability in cherry types and
3135 varieties for different characters.

3136 Differences among species and cultivars in plant response to mineral nutrient deficiency have been
3137 reported by many authors. Nutrient concentration and uptake by different plant cultivars are the
3138 most important criteria for identifying the existing genetic specificity of plant nutrition (Saric,
3139 1987). The tolerance in a given plant species or genotype to nutrient stress is closely related to its
3140 nutrient use efficiency. For a given genotype, nutrient use efficiency is reflected by the ability to
3141 produce a high yield in a soil that is limiting in one or more mineral nutrients for a standard
3142 genotype (Graham, 1984). The exploration of genotypic variability in the responses of sour cherry
3143 cultivars to mineral nutrient deficiency may allow identification of physiological or biochemical

3144 tools to screen tolerant varieties and to improve the productivity of this plant. Variability among
3145 cultivars in potential for uptake of mineral nutrients is caused by factors which are at least partly
3146 under genetic control. However, balanced nutrition is required for maximum tree performance and
3147 foliage is the tissue most generally used in diagnosis.

3148 Although there is extensive research on the pear, literature does not report enough data on the
3149 nutritional properties of pear based on this variety. The objective of this study was to determine
3150 mineral status of 29 sour cherry (*Prunus cerasus* L.) samples those were collected from Aegean
3151 Region of Turkey.

3152

3153 **Material and methods**

3154 A study was conducted in Aegean Region of Turkey for assessing the leaf nutrient contents of
3155 early-ripening sour cherry (*Prunus cerasus* L.) cultivars. Samples were collected as scion woody in
3156 August from an area between 38°45'-41°10' N latitude and 27°50'-37°05' E longitude. Elevation was
3157 ranged from 275 to 1150 m. A total of twenty-nine types were collected from wild and feral
3158 populations growing Aegean Region of Turkey. Trees have been marked for further visit and
3159 recollection. The samples were T-budded onto *Prunus mahaleb* seedlings in the nursery. Budded
3160 trees were planted in the experimental fields with 6x 6 m. spacing with tree replicates for each type.
3161 Routine cultural practices were performed. After three years of growth, plant leaf samples and fruit
3162 samples were collected and analyzed.

3163 Soil characteristics of experimental soil can be summarized as; sandy loam texture, slightly alkaline
3164 reaction, low CaCO₃, low organic matter, high exchangeable K and Ca contents, normal levels of N
3165 and plant available P, K, Ca, Mg, Fe, Zn, Mn and Cu. These values represent the typical Aegean
3166 region soil characteristics in Turkey.

3167 Leaf samples were taken on mid may, with five replications. Fruit samples were collected at
3168 maturity stage. Leaf and fruit samples were dried at 65 °C for 48 h and ground for determination of
3169 plant analysis. In dried plant samples, total N were determined by Kjeldahl method. Plant tissues
3170 were digested in aqua regia (1:3 HNO₃/HCl) for mineral analysis. In wet ashed plant samples total P
3171 were determined by molibdophosphoric yellow colour method, total K, Ca, Mg, Fe, Zn, Mn and Cu
3172 were determined by atomic absorption spectrophotometry (FAAS) under optimised measurement
3173 conditions.

3174 Statistical analyses were performed by using SPSS-16 for Windows program.

3175

3176 **Results and discussion**

3177 In this study, total quantities of plant nutrients accumulated within the leaf and fruits were used to
3178 make provisional estimates of uptake of macro and micro plant nutrients by sour cherry trees.

3179 Table 1 shows that the plant leaf mineral concentrations of sour cherry cultivars and statistic
3180 differences among the cultivars. All investigated nutrients were recorded significantly different
3181 within the cultivars. Plant leaf mineral contents of sour cherry cultivars were recorded in higher
3182 concentrations both in 1515 and 1530 cultivars for all elements. But leaf mineral contents of sour
3183 cherry cultivars were recorded in lower concentrations both in 1512 and 1519 cultivars for all
3184 elements.

3185 Leaf nitrogen concentrations of many cultivars were recorded in nitrogen deficiency level. The
3186 concentrations of all elements except Zn were recorded in normal plants nutrient limits. Only
3187 cultivar 1519 contained Zn concentration in acceptable range according to referenced limits.
3188 Slightly alkaline reaction of experimental soil could be a cause of zinc deficiency in evaluated
3189 cultivars. This also shows that the genotypic sensitivity of these sour cherry cultivars to zinc
3190 nutrition in slightly alkaline soil conditions and to nitrogen nutrition in low soil organic matter.
3191 According to these data, it is important to planning a balanced nitrogen and zinc fertilization
3192 programme for these cultivars in low soil organic matter and slightly alkaline soil conditions.

3193 Table 2 shows that the fruit mineral concentrations of sour cherry cultivars and statistic differences
3194 among the cultivars. As in leaf data, all investigated nutrients were recorded significantly different
3195 within the cultivars. Fruit mineral contents of sour cherry cultivars were recorded in higher
3196 concentrations both in 1507 and 1525 cultivars for all elements. But fruit mineral contents of sour
3197 cherry cultivars were recorded in lower concentrations both in 1512 and 1538 cultivars for all
3198 elements. Leaf mineral concentrations were recorded higher than that of fruit minerals. According
3199 to these data, genotype 1512 has contained the least mineral contents for all elements both in leaf
3200 and fruit tissues.

3201

3202 **Conclusion**

3203 In breeding programs priorities are generally given to traits as adaptability to environmental
3204 conditions, high yield and fruit quality or disease resistance. However, mineral nutrition may create
3205 marked effects on above mentioned characteristics. In this regard, significant differences were
3206 found among the sour cherry cultivars. Among the Sour cherry cultivars, the concentration of plant
3207 leaf nutrients in 1515 and 1530 cultivars were higher than the other investigated cultivars at the
3208 same soil conditions. A parallel study to this research conducted by authors also showed that these
3209 cultivars were taken the first place according to weighed-ranked method with regard to some
3210 characteristics such as yield, fruit size and earliness. These results could be helpful in development
3211 of reliable screening parameters for selection of high nutrient containing and high nutrient-use
3212 efficient cultivars in breeding programs.

3213

3214 **Acknowledgements**

3215 This study was supported by Akdeniz University. The authors wish to thank the Management of
3216 Vocational School of Technical Sciences and Aegean Agricultural Research Institute for their
3217 valuable contribution and technical support.

3218

3219 **References**

3220

3221 **Anonymous (A&L Canada Laboratories Inc.) (2002).** Cherry nutrition Deficiency symptoms. Fact
3222 sheet no: 13, Revised 11, London.

3223 **Bekefi Z., Apostol J. and Boronkay G. (2000).** Flowering dynamics in the Hungarian sour cherry
3224 genebank. Proceeding of the EUCAPRPIA Symposium on Fruit Breeding and Genetics, Dresden,
3225 Germany, 6-10 September.1999. (Eds. Geibel, M. And Fisher, M.) Vol 1, No:538,117-122.

3226 **Christensen J.V. (1978).** Evaluation of sour cherry cultivars. Saertry of Tidsskrift for Plant Cool.
3227 82:288-396.

3228 **Cordeiro-Rodrigues L., Ortiz-Marcide J.M., Morales-Corts M.R. and Socias-i-Company R.**
3229 **(2000).** Characterization of local varieties of sweet cherry and sour cherry from the Iberian
3230 Peninsula on the basis of morphology and biochemical componenets. Jornadas de experimentacion
3231 en futicultura, Zaragoza, Spain. ITEA. Volumen Extra No.21, 187-197.

3232 **Graham, R. D. (1984).** Breeding for nutritional characteristic in cereals. In '*Advances in Plant*
3233 *Nutrition*' (P. B. Tinker and A. Lauchli, eds.), Vol.1, pp. 57-102. Praeger, New York.

3234 **Saric, M. R. (1987).** Progress since the first international symposium : 'Genetic aspects of plant
3235 mineral nutrition'. Beogard, 1982, and perspectives of future research. *Plant and Soil.* 99: 197-209.

3236 **Vestrheim S. (1973).** Evaluation of sour cherry cultivars. Reports of the Agricultural University of
3237 Norway, Vol.52, (30):1-23.

3238

3239

3240

3241

3242

3243 Table 1. Leaf mineral element concentrations of 'sour cherry cultivars (*P. cerasus* L.) at 120 days
3244 after full bloom and the three year after grafting.

3245 Tablo 1. Vişne (*P. cerasus* L.) çeşitlerinin aşılama sonrası üçüncü yılında ve çiçeklenmeden 120
3246 gün sonra yapraklarında belirlenen mineral element konsantrasyonları

Cultivar No.	Cultivars	N	P	K	Ca	Mg	Fe	Zn	Mn	Cu
		%, dw					mg kg ⁻¹ , dw			
1	1507	2,21	0,34	1,84	1,90	0,65	50,4	15,8	62,4	9,6
2	1509	1,97	0,29	1,54	1,70	0,51	45,9	17,1	54,0	10,2
3	1510	2,12	0,35	1,55	1,74	0,52	41,0	15,5	58,4	8,4
4	1511	1,99	0,31	1,54	1,72	0,52	43,5	15,3	54,9	8,5
5	1512	2,60	0,37	2,05	2,25	0,67	56,7	21,0	71,4	12,8
6	1514	1,97	0,29	1,54	1,70	0,51	42,9	15,1	54,0	8,2
7	1515	1,84	0,25	1,44	1,60	0,43	40,3	14,2	47,7	7,7
8	1517	2,09	0,33	1,63	1,81	0,54	45,6	16,1	57,4	8,7
9	1518	2,48	0,35	1,86	2,06	0,62	51,9	18,3	68,4	9,9
10	1519	3,38	0,52	2,64	2,92	0,88	75,7	26,0	92,7	15,0
11	1520	1,90	0,28	1,40	1,69	0,49	40,3	14,6	52,0	7,5
12	1522	1,94	0,29	1,52	1,68	0,50	42,4	16,0	53,4	7,6
13	1523	2,06	0,33	1,63	1,91	0,52	48,2	15,0	55,7	9,2
14	1524	1,92	0,28	1,60	1,61	0,50	42,9	15,8	52,7	8,0
15	1525	2,26	0,33	1,77	1,85	0,59	46,3	17,4	62,0	9,4
16	1326	1,95	0,26	1,58	1,64	0,45	41,3	15,6	50,0	7,9
17	1528	1,94	0,29	1,52	1,68	0,53	42,4	16,0	58,4	8,1
18	1529	1,85	0,25	1,34	1,55	0,48	42,3	14,2	50,7	9,7
19	1530	1,82	0,27	1,43	1,68	0,45	37,8	12,7	48,0	7,6
20	1531	2,11	0,31	1,65	1,83	0,55	46,1	16,3	58,0	8,8
21	1532	1,80	0,27	1,44	1,65	0,45	40,3	13,2	52,7	6,7
22	1533	1,99	0,25	1,56	1,72	0,57	43,5	16,3	54,7	8,3
23	1535	2,24	0,33	1,65	1,83	0,58	46,8	17,2	61,4	9,3
24	1538	2,19	0,37	1,51	1,81	0,54	45,6	16,1	57,4	8,7
25	1539	1,87	0,28	1,71	1,58	0,49	43,8	14,0	50,0	7,6
26	1540	2,21	0,34	1,74	1,90	0,60	52,4	16,1	67,4	8,6
27	1541	1,97	0,29	1,58	1,70	0,55	46,9	15,1	55,0	8,2
28	1542	2,02	0,27	1,56	1,74	0,55	44,0	14,5	55,4	8,4
29	1543	1,99	0,29	2,03	1,82	0,56	41,5	16,3	54,5	8,1
Mean		2,09	0,31	1,65	1,81	0,50	42,6	16,1	52,4	9,7
Min-Max Values		1,82-	0,27-	1,43-	1,58-	0,47-	39,8-	14-26	50-92,7	7,6-14
St. Deviation		0,311	0,464	0,254	0,269	0,081	6,793	2,401	8,551	1,281
Sig. (2-tailed)		**	**	**	**	**	**	**	**	**
Mineral limits		2,2-3,4	0,16-	1-3	0,7-3	0,4-0,9	nls	25	nls	nls

3247 ** : Significant levels at 1 %; nls: no limitation set

3248 Table 2. Fruit mineral contents of Sour Cherry cultivars (*P. cerasus* L.) at maturity stage and the
 3249 three years after grafting.

3250 Tablo 2 Vişne (*P. cerasus* L.) çeşitlerinin aşılama sonrası üçüncü yılında olgunluk aşamasındaki
 3251 meyvelerinde belirlenen mineral element konsantrasyonları.

Cultivar No.	Cultivars	N	P	K	Ca	Mg	Fe	Zn	Mn	Cu
		% dw					mg kg ⁻¹ , dw			
1	1507	1,82	0,173	0,663	0,143	0,173	22,0	8,47	4,59	5,51
2	1509	1,35	0,158	0,640	0,125	0,163	20,4	8,00	4,36	5,40
3	1510	1,45	0,168	0,648	0,135	0,168	21,4	8,22	4,46	5,35
4	1511	1,16	0,139	0,533	0,115	0,139	17,7	6,81	3,69	4,43
5	1512	1,06	0,128	0,488	0,105	0,128	16,2	6,23	3,38	4,05
6	1514	1,34	0,162	0,618	0,133	0,162	20,5	7,89	4,28	5,13
7	1515	1,11	0,148	0,534	0,127	0,143	18,5	6,72	3,75	4,37
8	1517	1,17	0,141	0,540	0,116	0,173	17,9	6,89	3,74	4,48
9	1518	1,16	0,139	0,566	0,120	0,132	17,7	6,91	3,69	4,33
10	1519	1,51	0,182	0,696	0,150	0,182	23,1	8,88	4,82	5,78
11	1520	1,17	0,138	0,520	0,113	0,138	17,5	6,72	3,65	4,37
12	1522	1,09	0,133	0,498	0,116	0,121	16,7	6,27	3,32	4,05
13	1523	1,17	0,140	0,549	0,127	0,136	18,0	7,04	3,77	4,64
14	1524	1,38	0,167	0,637	0,137	0,167	21,2	8,13	4,41	5,29
15	1525	1,96	0,236	0,904	0,195	0,236	30,0	11,54	6,26	7,51
16	1326	1,10	0,133	0,507	0,114	0,135	16,5	6,45	3,41	4,11
17	1528	1,13	0,136	0,520	0,112	0,136	17,3	6,64	3,60	4,32
18	1529	1,28	0,155	0,592	0,127	0,155	19,7	7,55	4,10	4,91
19	1530	1,11	0,134	0,514	0,111	0,134	17,6	6,66	3,66	4,27
20	1531	1,31	0,158	0,605	0,120	0,158	20,1	7,94	4,19	5,02
21	1532	1,10	0,141	0,517	0,105	0,132	17,1	6,49	3,61	4,31
22	1533	1,13	0,136	0,520	0,112	0,136	17,3	6,64	3,60	4,32
23	1535	1,05	0,125	0,490	0,106	0,137	16,4	6,35	3,42	4,10
24	1538	1,06	0,128	0,488	0,105	0,128	16,2	6,23	3,38	4,05
25	1539	1,23	0,148	0,566	0,122	0,148	18,8	7,22	3,92	4,70
26	1540	1,07	0,129	0,494	0,106	0,129	16,4	6,31	3,42	4,10
27	1541	1,16	0,130	0,520	0,110	0,146	17,4	6,71	3,79	4,53
28	1542	1,30	0,156	0,598	0,129	0,156	19,9	7,64	4,14	4,97
29	1543	1,25	0,152	0,559	0,120	0,156	18,6	7,14	3,87	4,64
Mean		1,23	0,149	0,569	0,122	0,150	18,9	7,27	3,94	4,73
Min-Max Values		1,05-	0,125-	0,488-	0,105-	0,121-	16,2-	6,23-	3,32-	4,05-
St. Deviation		0,188	0,227	0,087	0,189	0,023	2,889	1,111	0,603	0,724
Sig. (2-tailed)		**	**	**	**	**	**	**	**	**

3252 **: Significant levels at 1 %.

Characterization and Evaluation of Sour Cherry (*Prunus cerasus* L.) Genetic Resources in Aegean Region of Turkey

M. KUBİLAY ÖNAL, BÜLENT TOPCUOĞLU

AkdenizUniversity, Vocational High School of Technical Sciences

Kampus 07058 Antalya, Turkey

*Corresponding author: konal@akdeniz.edu.tr

Abstract

The purpose of this study was to determine the phenological, pomological and technological characters of 29 sour cherry (*P. cerasus* L.) types collected from Aegean Region of Turkey. The types were evaluated for the characters of first blossoming, full bloom, end of blossoming, harvesting time, yield, fruit weight, fruit flesh/seed, juice content, total soluble solids/acidity, pH, juice colour, fruit shape, skin colour, aroma and attractiveness. It was found that the earliest blooming typ was 1528, which usually bloom at the end of March. Harvesting time began in the beginning of June. The earliest ripening typ was 1528. The types that ripen late were 1533 and 1539 about end of June. Average fruit weight was around 4.0 g-5.0g. Four different colours were observed for fruit skin. The colour of fruit skin ranges from cherry, dark cherry, dark red to light cherry. There was certain amount of variability among sour cherry types which were collected from Aegean Region.

Keywords: evaluation, sour cherry, genetic resources

Türkiye'nin Ege Bölgesi Vişne (*Prunus cerasus* L.) Genetik Kaynaklarının Karakterizasyonu ve Değerlendirilmesi

Özet

Çalışmada; Ege Bölgesi'nin değişik yörelerinden toplanan 29 vişne (*P. cerasus* L.) tipinin değerlendirilmesi ve karakterizasyonu yapılmıştır. Tiplerin; fenolojik özellikleri (ilk çiçeklenme, tam çiçeklenme, çiçeklenme sonu ve hasat tarihi), pomolojik özellikleri (meyve/ çekirdek oranı, meyve suyu içeriği, suda çözünebilir kurumadde/asit oranı, pH, meyve suyu rengi, meyve şekli,

3286 meyve rengi, aroma ve albeni) ve verim deęerleri belirlenmiřtir. Tipler arasında geniř bir
3287 varyasyonun olduęu gzlenmiřtir.

3288

3289 **Anahtar kelimeler:**viřne, genetic kaynakları, deęerlendirme.

3290

3291

3292 **Introduction**

3293 The 'Ktahya' sour cherry (*Prunus cerasus* L.) is the most planted cultivar in Turkey commercial
3294 orchards. This cultivar is self-compatible and characterized by regular and high yield. The fruit is of
3295 "morello" type, medium to big in size, with dark red and thin skin. The flesh is red, medium-firm,
3296 juicy, quite sour, aromatic and of high quality. (zaęiran, 1977a; zaęiran, 1977b; z, 1988;
3297 Burak et.al.,1999). The numerous positive traits of this cultivar should make it interesting for
3298 plantation in other countries.

3299 Anatolia is located in two of eight plant origin centers suggested by Vavilov (1951), "Near Eastern
3300 Center" and "Mediterranean Center". Several species of fruit trees or their wild relatives are
3301 indigenous to Turkey like fig, walnut, almond, apple, pear, plum, sour cherry and olive (Sykes
3302 1972).

3303 This species is indigenous to Turkey and there is a wide range variation in fruit characteristics. Sour
3304 cherry is widely planted in the Central Anatolia and Aegean Region of Turkey. These districts have
3305 the best climatic factors, such as brilliant illumination, low temperature and great difference in
3306 temperature between day and night. A few studies (Vesterheim 1973; Christersen 1978; Bekefi *et*
3307 *al.* 2000; Cordeir Rodrigues *et al.* 2000) have revealed wide variability in cherry types and varieties
3308 for different characters.

3309 Thirty-four sour cherry varieties were tested for fruit size and weight, flesh/kernel ratio, percentage
3310 juice yield, juice colour, acid content and an evaluation of the quality of the processed product
3311 (Liverani *et al.*1981). Moltalti and Camorani (1982) evaluated for dates of flowering and maturity,
3312 habit, peduncle length, percentage of soluble dry matter, pH of the juice, acid content of the juice,
3313 fruit size, fruit weight, fruit shape, colour of the skin, flesh and juice, and flesh consistency of 21
3314 cultivars.

3315 In a trial with 6 varieties during 1986-90 at Zajecar on the Beli Timok river in Serbia, Oblacinska
3316 was the earliest to flower, and mean fruit weight ranged from 3,1 g. in Oblacinska to 5,8 g. in
3317 Hajmanova Konzervna. Total sugar content of fruit varied from 8,5 % in Oblacinska to 9,7 % in
3318 Hajmanova Konzervna. Solible solids was highest (at 15,2 %) in Cacanski Rubin (Miletic,1991).

3319 Eight varieties and three selection types were determined from Hungarian breedin programme for
3320 different characters (Apostol *et al.*1993)

3321 Six local types and three foreign varieties were examined in Southeast Turkey. Two local types
3322 were found to be more suitable for that region (Karaca *et al.* 1998). Morphological variation among
3323 78 local races of sour cherry provided from the Nordic Gene Bank's were observed in Palkane,
3324 Southwestern Finland (Palonen *et al.* 2000). On the basis of fruit characteristics, 32 sour cherry
3325 samples were classified as morellos, 40 samples as ameralles, and 5 samples being distinct from the
3326 other amarells were proposed to be called Rymattyla-type cherries.

3327 The beginning, peak and end of flowering periods of 90 sour cherry evaluated. The beginning and
3328 duration of flowering of the evaluated cultivars differed in the individual years of studies depending
3329 on the weather conditions. The earliest flowering was noticed at the end of the second 10 days period
3330 of April, while the latest was on the first day of May. Depending on the year, the earliest flowering
3331 cultivars flowered 6 to 12 days earlier than the latest flowering (Hodun and Hodun 2002).

3332 The complete characterization as well as the conservation of autochthonous cultivars is considered
3333 of great importance in order to avoid the loss of their germplasm (Rodrigues *et al.*, 2008).

3334 Forty one genotypes of the 'Oblačinska' sour cherry collection, to determine the correlation among
3335 the traits, to identify the most useful variables for discrimination among the genotypes and to detect
3336 relationships between the genotypes (Rakonjac *et al.*, 2010).

3337 The objective of this study was to determine phenological, pomological and technological
3338 characters of 29 sour cherry (*P. cerasus* L.) samples collected from Aegean Region of Turkey.

3339

3340 **Material and Methods**

3341 Samples were collected as scion woody in August from an area between 38°45'-41°10' N latitude
3342 and 27°50'-32°05' E longitude. Elevation ranged from 225 to 950 m. A total of twenty-nine types
3343 was collected from wild and feral populations growing Aegean Region of Turkey. Trees have been
3344 marked for further visit and recollection.

3345 The samples were T-budded onto *Prunus mahaleb* seedlings in the nursery. Budded trees were
3346 planted in the experimental fields with 6x 6 m. spacing with tree replicates for each type. Routine
3347 cultural practices were performed.

3348 Types were evaluated for important phenological, pomological and technological characteristics in
3349 order to determine their importance for horticulture and breeding during the period of three years.
3350 First blossoming, full bloom, end of the blossoming and harvesting time were recorded. Flowering
3351 time were observed three times a week. Some other characteristics such as fruit shape, skin colour,
3352 pH value of juice, aroma, fruit flesh/seed ratio were also recorded.

3353

3354 **Results and Discussion**

3355 All types evaluated for phenological characteristics are listed in Table 1. It was found that the
3356 earliest blooming typ was 1528, which usually bloom at the end of March. First blossoming date of
3357 types were between 25 March and 17 April. The period from the begining of blooming to the end of
3358 blooming was 20 days, changing from years to years.

3359 Harvesting time began in the beginning of June. The earliest ripening typ was 1528. The types that
3360 ripen late were 1524 and 1515 about end of June. Harvesting period was approximately 20 days.

3361 Several pomological characters are given Table 2 and 3. Fruit weight of the accessions ranged
3362 between 3.3 g and 5.9 g, sample 1514 the highest and sample 1538 the lowest. Average fruit weight
3363 was around 4.0 g.-5.0g 65% of types were placed between the variation limits of 4.0 to 5.0 g in fruit
3364 weight. From the point of view of average fruit weight, the types collected were generally in a good
3365 position as in similar researchs carried out both in Turkey and in the other countries. For example
3366 Karaca *et al.* (1998) 3.5-4.0 g and Miletic (1991) recorded 3.1.-5.8 g. On the contrary, Nikolic *et al.*
3367 (1991) cited that Gaga variety had been determined as having the largest fruit weight with more
3368 than 8.0 g in Serbian.

3369 Fruit flesh/seed ratio values calculated were between 6.9 in type 1525 and 4.6 in types 1522, 1526
3370 and 1514. The total soluble solids (TSS)/acidity of types varied from 7.5 to 13.9. Type 1525 had
3371 the highest followed by type 1519 with 10.7, and type 1507 had the lowest TTS/acidity value with
3372 7.5.

3373 A large variation was noticed among types for the rate of fruit juice, type 1535 had the lowest,
3374 66.3%, and type 1510 had the highest, 84%. The analysis showed that pH values changed from 2.8
3375 (type 1520) to 3.3 (types 1528 and 1329). It was found out that there was no big differences among
3376 types for pH values.

3377 The fruit yield per tree showing considerably large variation was recorded from a single harvest. It
3378 ranged from a minimum of 3.8 kg to a maximum of 20.3 kg. The highest yields were obtained from
3379 type 1539 and 1532 as 20.3 and 19.8 kg/tree, respectively. Type 1522 produced the lowest yield as
3380 3.8 kg/tree. Yield correlated not strongly but significantly and negatively with fruit weight.

3381 Four different fruit shapes were determined. Round shape samples were more common than the
3382 others with approximately 60% of all types. Flat-round and heart shape types being approximate in
3383 numbers were much more than kidney shape (only one sample).

3384 Four different colours were observed for fruit skin. The colour of fruit skin ranges from cherry, dark
3385 cherry, dark red to light cherry.

3386 There were five different colours for fruit juice, cherry, dark cherry, dark red, red, light cherry.
3387 Cherry and dark cherry colours together were more common than the others for fruit skin colour
3388 and fruit juice colour making up a total of 29 and 14 of all types.

3389 Type 1515 had very good aroma and they were found to be suitable for fresh consumptions. Poor
3390 aroma was observed for more than half of the sour cherry types studied.

3391 Attractiveness was scored as very good, good, average and poor. Only one type had a very good
3392 appraisal. Around 85 percent of types had good and average attractiveness. Close relationships were
3393 observed between attractiveness and fruit weight and fruit skin colour.

3394 We have also conducted an evaluation of economically valuable characteristics in order to identify
3395 the genotypes with a good potential for cultivation or for exploitation in breeding programs.

3396

3397 **References**

3398

3399 **Apostol J., Gyuro F., Gergely I. and Cossio F. 1993.** Genetic basis for breeding sour cherry in
3400 Hungary. *Indirizzi nel miglioramento genetico e nella coltura del ciliegio, atti del convegno,*
3401 *Verona, 21-22 giugno, 51-60.*

3402 **Bekefi Z., Apostol J. and Boronkay G. 2000.** Flowering dynamics in the Hungarian sour cherry
3403 genebank. *Proceeding of the EUCAPRPIA Symposium on Fruit Breeding and Genetics, Dresden,*
3404 *Germany, 6-10 September. 1999. (Eds. Geibel, M. And Fisher, M.) Vol 1, No:538, 117-122.*

3405 **Budan S. and Stoian I. 1996.** Genetic resources in the Romanian sour cherry breeding program.
3406 *Proceedings of the international cherry symposium, Budapest, Hungary, 14-18 June. Acta-*
3407 *Horticulturae. No.410, 81-86.*

3408 **Burak, M., Erbil, Y. and Kaynaş, K. 1999.** Adaptation of sour cherry cultivars. *Atatürk Central*
3409 *Horticultural Research Institute. Publication no:126. Yalova.*

3410 **Christensen J.V. 1978.** Evaluation of sour cherry cultivars. *Saertry of Tidsskrift for Plant Cool.*
3411 *82:288-396.*

3412 **Christensen J.V. 1990.** A review of an evaluation of 95 cultivars of sour cherry. *Tidsskrift-for-*
3413 *Planteavl. 94:1, 51-63.*

3414 **Cociu V. 1994.** Genetic resources of prunus species in Romania. *Plant Genetic Resources*
3415 *Newsletter. No.100, 22-23.*

3416 **Cordeiro-Rodrigues L., Ortiz-Marcide J.M., Morales-Corts M.R. and Socias-i-Company R.**
3417 **2000.** Characterization of local varieties of sweet cherry and sour cherry from the Iberian Peninsula
3418 on the basis of morphology and biochemical components. *Jornadas de experimentacion en*
3419 *futicultura, Zaragoza, Spain. ITEA. Volumen Extra No.21, 187-197.*

- 3420 **Fogle H.W. 1975.** Cherries. In: Janick, J., Moore, J.N., eds. Advances in Fruite Breeding. Purdue
3421 University Press, USA.
- 3422 **Hodun M. and Hodun G. 2002.** Evaluation of flowering of 90 sour cherry cultivars and their
3423 classification in regard to the season of blooming. Annales-Universitatis-Mariae-Sklodowska.
3424 Sectio-EEE,- Horticultura. 10:201-206.
- 3425 **Karaca R., Akkök F. and Atlı H.S.1995.** Adaptation of sour cherry cultivars. Second National
3426 Congres of Horticulture, Adana, Turkey, 3-6 October 1995, Volum I, pp:243-247.
- 3427 **Liverani A., Cobianchi D. and Bergamini A. 1981.** Studies on the suitability of sour cherry
3428 varieties for industrial processing. Informatore-Agrario. 37:50, 18459-18463.
- 3429 **Miletic R. 1991.** Pomological and technological characteristics of the fruit in some sour cherry
3430 varieties grown in the Timok border region. Jugoslovensko-Vocarstvo. 25:1-2, 39-43.
- 3431 **Montalti P. and Camorani F. 1982.** Contribution to the knowledge and evaluation of sour cherry
3432 cultivars. Informatore-Agrario. 38:13, 20235-20241.
- 3433 **Nikolic M.; Cerovic,R., Rankovic,M. and Ogasanovic, D. 1991.** Characteristics of selected local
3434 types of sweet cherry and sour cherry. Jogoslovensko-Vocarstvo. 25:1-2,27-31.
- 3435 Nikolic,D.,Rakonjac, V., Milatovic, D. and Fotiric, M. 2010. Multivariate analysis of vineyard
3436 peach [*Prunus persica* (L.) Batsch.] germplasm collection.Euphytica, 171, pp. 227–234.
- 3437 **Öz, F. 1988.** Cherry. TAV Publications. Publication no:16. 72 p.,Yalova.
- 3438 **Özçağiran, R. 1977a.** Cherry. Ege University Faculty of Agriculture publication. Publication
3439 no:328, 106p., Bornova, İzmir.
- 3440 **Özçağiran, R. 1977b.** Cherry. Pomological characters of some important sour cherry varieties.
3441 Journal of Agricultural Faculty of Ege University, 14(3):311-328. Bornova, İzmir.
- 3442 **Palonen P., Uosukainen M., Aaltonen M., Laurinen E. and Kaufmane, E. 2000.** Local races of
3443 sour cherry and plum in the Nordic gene Bank's Prunus clone archive in Finland. Proceeding of the
3444 International Conference Fruit Production and fruit Breeding, Tartu, Estonia, 12-13 September,
3445 138-142.
- 3446 **Papic V. 1987.** Pomological and biochemical features of the fruit in some sour cherry varieties.
3447 Jugoslovensko-Vocarstvo. 21:4, 17-21.
- 3448 **Perez, S., Montes, S. and Mejia, C. 1993.** Analysis of peach germplasm in Mexico
3449 J. Am. Soc. Hortic. Sci., 118, pp. 519–524.
- 3450 **Rodrigues.,Morales, M.R., Fernandes, A.J.B. and Ortiz, J.M. 2008.** Morphological
3451 characterization of sweet and sour cherry cultivars in a germplasm bank at Portugal
3452 Genet. Resour. Crop Evol., 55, pp. 593–601.
- 3453 **Vera Rakonjac, V., Aksic, M.F., Nikolić, D., Milatović, D. And Colic, S. 2010.**

3454 Morphological characterization of 'Oblačinska' sour cherry by multivariate analysis. Scientia
3455 Horticulturae. Volume 125, Issue 4, 26 July 2010, Pages 679–684

3456 **Sykes J.T. 1972.** Propagation and collection techniques for fruit germplasm. Plant Propagator
3457 18:15-19.

3458 **Vavilov N.I. 1951.** The origin, variation, immunity, breeding of cultivated plant. Chron. Bot. 13:1-
3459 364.

3460 **Vestrheim S. 1973.** Evaluation of sour cherry cultivars. Reports of the Agricultural University of
3461 Norway, Vol.52, (30):1-23.

3462

3463

3464

3465

3466

3467

3468

3469

3470

3471

3472

interim

3473 Table 1. Some phenological characteristics of sour cherr types.

3474 Çizelge1. Vişne tiplerinde incelenen bazı fenolojik özellikler

Type no	Firs blossoming	Full bloom	End of blossoming	Harvesting time
1507	3.4-12.4	15.4-18.4	25.4-5.5	17.6-25.6
1509	4.4-11.4	15.4-18.4	19.4-7.5	17.6-25.6
1510	7.4-17.4	17.4-22.4	27.4-8.5	25.6-2.7
1511	4.4-12.4	15.4-19.4	26.4-5.5	23.6-25.6
1512	3.4-13.4	15.4-17.4	24.4-30.4	17.6-23.6
1514	4.4-17.4	15.4-24.4	26.4-4.5	17.6-30.6
1515	3.4-14.4	15.4-20.4	26.4-4.5	13.6-23.6
1517	2.4-13.4	16.4-17.4	26.4-2.5	13.6-17.6
1518	3.4-10.4	10.4-15.4	25.4-30.4	13.6-18.6
1519	3.4-13.4	15.4-20.4	27.4-6.5	17.6-23.6
1520	4.4-11.4	15.4-17.4	23.4-1.5	17.6-23.6
1522	4.4-13.4	15.4-17.4	22.4-4.5	17.6-23.6
1523	4.4-13.4	15.4-18.4	24.4-4.5	17.6-23.6
1524	3.4-13.4	15.4-20.4	25.4-4.5	17.6-23.6
1525	12.4-15.4	18.4-22.4	29.4-5.5	17.6-2.7
1326	4.4-13.4	16.4-18.4	22.4-1.5	17.6-23.6
1528	25.3-10.4	5.4-13.4	15.4-20.4	4.6-23.6
1529	3.4-13.4	15.4-19.4	23.4-6.5	13.6-23.6
1530	4.4-13.4	15.4-17.4	22.4-4.5	17.6-23.6
1531	4.4-13.4	15.4-18.4	24.4-4.5	17.6-23.6
1532	3.4-13.4	15.4-20.4	25.4-4.5	17.6-23.6
1533	12.4-15.4	18.4-22.4	29.4-5.5	17.6-2.7
1535	4.4-13.4	16.4-18.4	22.4-1.5	17.6-23.6
1538	5.4-10.4	15.4-18.4	23.4-3.5	17.6-18.6
1539	5.4-12.4	14.4-18.4	22.4-3.5	17.6-23.6
1540	3.4-15.4	15.4-20.4	25.4-5.5	13.6-23.6
1541	4.4-13.4	15.4-18.4	25.4-4.5	17.6-23.6
1542	3.4-13.4	15.4-20.4	27.4-6.5	17.6-23.6
1543	4.4-11.4	15.4-17.4	23.4-1.5	17.6-23.6

3475

3476

3477

3478 Table 2. Some pomological characteristics of sour cherry types.
 3479 Çizelge2. Vişne tiplerinde incelenen bazı meyvesel özellikler
 3480

Type no	Fruit shape	Skin color	Juice color	Aroma	Attractiveness
1507	Round	Chery	Dark cherry	Poor	Good
1509	Heart	Dark cherry	Cherry	Poor	Good
1510	Round	Dark cherry	Cherry	Good	Average
1511	Round	Cherry	Cherry	Average	Average
1512	Flat-round	Dark red	Dark cherry	Poor	Average
1514	Heart	Dark cherry	Dark cherry	Poor	Average
1515	Flat-round	Dark cherry	Dark cherry	Very good	Good
1517	Round	Light cherry	Cherry	Poor	Good
1518	Flat-round	Cherry	Dark cherry	Good	Good
1519	Round	Dark red	Dark cherry	Poor	Average
1520	Round	Dark red	Red	Poor	Good
1522	Flat-round	Dark red	Dark red	Poor	Good
1523	Round	Cherry	Dark red	Poor	Good
1524	Round	Cherry	Dark cherry	Poor	Good
1525	Round	Cherry	Dark cherry	Poor	Poor
1326	Round	Light cherry	Cherry	Poor	Good
1528	Flat-round	Cherry	Dark cherry	Poor	Good
1529	Round	Dark red	Red	Poor	Good
1530	Flat-round	Dark red	Dark red	Poor	Good
1531	Round	Cherry	Dark red	Poor	Good
1532	Round	Cherry	Dark cherry	Poor	Good
1533	Round	Cherry	Dark cherry	Poor	Poor
1535	Round	Light cherry	Cherry	Poor	Good
1538	Heart	Dark cherry	Dark cherry	Good	Average
1539	Kidney	Light cherry	Cherry	Poor	Very good
1540	Round	Light cherry	Dark red	Poor	Good
1541	Heart	Dark red	Cherry	Average	Poor
1542	Round	Dark red	Dark cherry	Poor	Average
1543	Round	Dark red	Red	Poor	Good

3481
 3482
 3483
 3484
 3485
 3486
 3487
 3488
 3489
 3490
 3491
 3492
 3493
 3494
 3495
 3496

3497 Table 3. Some pomological characteristics of sour cherry types.
 3498 Çizelge3. Vişne tiplerinde incelenen bazı meyvesel özellikler.
 3499

Type no	Fruit weight (g)	Fruit flesh/seed	Rate of juice (%)	Total soluble solids / acidity	pH	Yield (kg/tree)
1507	3,4	5,1	75,3	7,6	3,0	17,9
1509	4,2	5,2	74,6	8,2	3,0	16,8
1510	5,1	5,9	78,0	9,2	2,8	17,1
1511	5,0	5,3	73,0	8,6	2,9	9,6
1512	4,1	5,5	72,3	10,2	3,0	8,7
1514	4,2	5,8	73,0	9,9	3,1	9,5
1515	4,3	5,5	72,6	9,9	3,1	6,8
1517	3,9	5,2	70,5	8,2	3,0	11,4
1518	4,2	5,8	73,0	7,5	3,0	12,6
1519	4,7	6,3	75,0	10,7	3,0	7,6
1520	3,7	5,0	76,0	8,1	3,0	10,7
1522	4,8	5,3	75,0	7,6	2,9	3,8
1523	4,0	5,3	73,6	8,6	3,0	12,4
1524	4,4	5,7	67,3	9,8	3,1	9,8
1525	4,6	6,9	62,0	13,9	3,2	9,1
1326	4,1	5,4	76,0	7,8	3,0	8,5
1528	4,6	5,7	72,6	8,0	3,3	16,3
1529	4,1	5,9	73,3	9,1	3,3	11,2
1530	4,8	5,3	75,0	7,6	2,9	4,3
1531	4,0	5,3	73,6	8,6	3,0	12,4
1532	4,4	5,7	67,3	9,8	3,1	19,8
1533	4,6	6,9	62,0	13,9	3,2	7,4
1535	4,1	5,4	76,0	7,8	3,0	8,5
1538	5,9	6,9	65,0	9,5	3,0	6,8
1539	4,2	5,7	84,0	8,1	3,0	20,3
1540	5,5	5,7	73,3	8,3	3,0	17,2
1541	5,1	5,8	71,5	8,2	3,2	14,2
1542	4,7	6,3	75,0	10,7	3,0	7,6
1543	3,7	5,0	76,0	8,1	3,0	11,9
Mean	4,427	5,628	72,82	9,086	3,038	11,382
St Dev.	0,550	0,535	4,575	1,642	0,115	4,462
Sig.	**	**	**	**	**	**

3500
3501

The Clonal Selection Breeding of Akça Pear (*Pyrus communis* L.) in Aegean Region of Turkey

M. KUBILAY ÖNAL, BÜLENT TOPCUOĞLU

Akdeniz University, Vocational High School of Technical Sciences, Kampus 07058 Antalya, Turkey

*Corresponding author: konal@akdeniz.edu.tr

Abstract

Key words: Pear, Phenological and pomological observations, 'weighed-ranked' method.

Türkiye'nin Ege Bölgesi Akça Armut (*Pyrus communis* L.) Tiplerinin Kolonal Seleksiyonu

Özet

Anahtar kelimeler: Armut, fenolojik ve pomolojik gözlemler, tartılı derecelendirme.

3535 **Introduction**

3536 The pear originated in prehistoric times as a fruit crop. Cultivar development has been continuous
3537 since early days and a high level of improvement has been achieved. Pears are now grown in all
3538 temperate regions of the world (Layne and Quamme,1975).

3539 Species of pear belong to the genus *Pyrus*, the sub-family Pomoideae and the family Rosaceae. The
3540 genus *Pyrus* is believed to have arisen during the Tertiary period in the mountainous regions of
3541 Western China and to have evolved as it spread along the mountain chains to the east and west
3542 (Rubzov, 1944). Three centers of diversity for the genus have been identified by Vavilov (1951):
3543 the Chinese, the Central Asian and the Near Eastern / Asia Minor Centers. The Near Eastern Center
3544 is of special importance because it is believed that the domesticated forms of european pear (*Pyrus*
3545 *communis* L.) form which the modern cultivars are derived originated there.

3546 During long time course, many local pear cultivars that show different degrees of improvement or
3547 adaptability developed in Turkey (Browicz, 1972; Özbek, 1947 and 1978). Akça (*Pyrus communis*
3548 L.) is the earliest ripening pear cultivar in Turkey and has several different types (Ergun, 1989).

3549 In this research, the aim was to determine different types of Akça pear by surveying and select
3550 candidate types having high-yielding capacity and better fruit characteristics and adaptable to the
3551 conditions of Aegean Region of Turkey.

3552

3553 **Material and methods**

3554 The clonal selection studies were conducted in the villages and counties of Izmir, Aydın and
3555 Manisa provinces of Aegean region of Turkey on early-ripening Akça pear cultivar (*Pyrus*
3556 *communis* L.). As a result of selection studies 28 types were selected in the first step. In the second
3557 step, 12 out of them were found as promising ones by pre-evaluation studies and were taken into an
3558 adaptation trial at the Aegean Agricultural Research Institute (**AARI**) located near Menemen
3559 county of İzmir province.

3560 Budwoods were taken from each of original trees and budded on Quince A (**QA**) rootstocks. The
3561 experiment was established with 7 budded trees of each type of Akça cultivar at 6x6 spacing in a
3562 randomized block design. Akça cultivar has been already planted in orchard of Aegean Agricultural
3563 Research Institute (**AARI**) was used as a control in the experiment.

3564 For evaluation of those types, phenological and pomological characteristics were observed and
3565 yield efficiency (**productivity**) values were scored.

3566 Bud burst, first blossoming, full bloom, last blossoming and harvesting date were used as criteria
3567 for phenological observations.

3568 The shape, fruit weight, length of fruit stalk, taste, juiciness, texture and firmness of fruit flesh,
3569 ground and overcolour of fruit skin, russet amount in fruit skin, skin thickness, Total Soluble Solids

3570 (TSS) (%), aroma and eating quality of fruit were characteristics used for pomological evaluation
3571 (UPOV, 1974; IBPGR, 1983) and were scored as an average of 20 fruit samples taken 7 trees of
3572 each type, randomizely. For determination of colours, Methuen handbook of colour was used
3573 (Kornerup and Wanscher, 1978).

3574 Last year of the experiment, morphological characteristics of trees were observed and trees were
3575 classified as upright, semi-upright and spreading for tree habit and weak, intermediate and vigorous
3576 for tree vigor.

3577 Yield efficiency = productivity (kg / cm²) were obtained by dividing cumulative yield of each tree
3578 to 1 cm² of trunk cross-sectional area.

3579 'Weighed-ranked' method was employed in evaluating values (Büyükyılmaz et al.,1992).

3580 The characteristics such as yield, fruit size and earliness and their relative points, class values and
3581 class points were used as criteria in evaluation of types. Total points of types were obtained by
3582 multiplying relative points with class points of characteristics.

3583

3584 **Result and discussion**

3585 According to average of four-year phenological observations, dates for bud burst, first blossoming,
3586 full bloom and last blossoming were determined as 10-17/3, 23-29/3, 3-9/4, and 9-19/4,
3587 respectively. It was observed that harvesting dates of types varied between 13/6-7/7 (table 1).

3588 It was observed that there was no big difference for fruit shape among types. Fruit characteristics
3589 such as fruit width, fruit length, fruit stalk length and fruit flesh firmness were determined as 35.7-
3590 54.1 mm, 51.8-71.2 mm 24.7-48.5 mm and 9-12 lb/cm² respectively (Table 2. Fruit characteristics
3591 such as fruit weight and total soluble solids were determined as 29.4-90.2g and 11-15 %, also
3592 (Table 3).

3593 There was no difference for characteristics such as taste, juiciness, texture and firmness of fruit
3594 flesh, ground and overcolour of fruit skin, russet amount in fruit skin, skin thickness and aroma and
3595 eating quality of fruit among types.

3596 Cumulative yield varied between 9.112-78.912kg/tree. Yield efficiency (productivity) was found as
3597 0.077-0.350 kg/cm² among types.

3598 The ranking of types in respect, to points obtained by 'weighed-ranked ' method is given at table 3.

3599 The types having high points at the table, viz. nos 2036, 2051 and 2103, were chosen as promising
3600 types suitable to the conditions of Aegean Region of Turkey.

3601

3602 **References**

3603

3604 **Brovicz, K. 1972.** *Pyrus* L. In: Flora of Turkey and The East Aegean Islands. Ed.P.H.Davis, Vol. 4.
3605 pp. 160-169. Edinburgh University Press.

3606 **Büyükyılmaz, M., Bulagay, A.N. and Burak, M. 1992.** Report of Clonal Selection on Akça and
3607 Ankara Pear Cultivars (**in Turkish**) (Marmara Region of Turkey), Yalova, Turkey.

3608 **Ergun, M. 1989.** A Report of Marketing Problems of Pear Production (**in Turkish**) Yalova,
3609 Turkey.

3610 **IBPGR. 1983.** Descriptor List for Pear (*Pyrus*). AGPG: IBPGR / 82 / 57.

3611 **Kornerup, A. and Wanscher, J.H. 1978.** Methuen Handbook of Colour. London, UK.

3612 **Layne, R.E.C. and Quamme, H.A. 1975.** Pears. In: " Advances in Fruit Breeding", eds. J.Janick
3613 and J. N. Moore, pp. 38-70. Purdue University Press, West Lafayette, Indiana, USA.

3614 **Özbek, S. 1947.** Pear Culture in Turkey and Important Turkish Pear Cultivars (**in Turkish**).
3615 Agriculture Institute Publication.

3616 **Özbek, S. 1978.** Special Fruit Production (**in Turkish**). Agriculture Faculty of Çukurova
3617 University Publ. No. 128.

3618 **Rubzov, G.A. 1944.** Geographical distribution of the genus *Pyrus* and trends and factors in its
3619 evolution. Amer. Nat. 78: 358-366.

3620 **UPOV. 1974.** Guidelines for the Conduct of Tests for Distinctness, Homogeneity and Stability.
3621 Pears. UPOV / TG / 15 / 1.

3622 **Vavilov, N.I. 1951.** The Origin, Variation, Immunity and Breeding of Cultivated Plants. Ronald
3623 Press, NewYork, USA.

3624

3625 Table 1. Some phenological characteristics of Akça pear.

3626 Çizelge 1. Armut tiplerinde incelenen bazı fenolojik özellikler.

Type no	Bud burst	Firs blossoming	Full bloom	End of blossoming	Harvesting time
2036	10-14.03	25-29.03	05-09.04	13-17	20-25.06
2039	12-16.03	23-29.03	03-07.04	11-15	20-25.06
2042	10-15.03	25-27.03	05-07.04	13-19	20-25.06
2043	11-16.03	23-29.03	03-09.04	09-15	13-17.06
2045	12-17.03	27-29.03	07-09.04	17-19	29.06-02.07
2051	10-17.03	25-29.03	05-07.04	15-19	29.06-02.07
2099	13-17.03	23-29.03	03-07.04	13-15	20-25.06
2100	14-17.03	23-27.03	03-05.04	09-13	13-17.06
2101	11-14.03	25-27.03	05-09.04	15-19	20-25.06
2102	11-15.03	27-29.03	07-09.04	17-19	29.06-02.07
2103	13-16.03	25-27.03	07-09.04	17-19	07-09.07
2104	12-15.03	23-25.03	03-05.04	09-13	13-17.06
Akça (C)	14-17.03	25-29.03	07-09.04	15-19	29.06-02.07

3627 C. Control

3628

3629

3630

3631 Table 2. Some phenological characteristics of Akça pear.

3632 Çizelge 2. Armut tiplerinde incelenen bazı pomolocik özellikler.

Type no	Fruit width	Fruit length	Fruit stalk length	fruit flesh firmness
2036	48.2-54.1	63.7-70.6	29.3-34.5	9.2-11.4
2039	41.6-48.1	62.3-67.4	41.7-46.8	9.7-10.8
2042	39.3-44.2	53.6-61.4	35.7-46.2	10.3-11.7
2043	35.7-42.8	52.4-60.6	29.6-38.5	10.7-12.0
2045	44.3-51.4	55.2-62.7	38.9-47.6	9.6-10.9
2051	47.8-53.7	64.3-71.2	25.4-37.8	10.3-10.7
2099	46.2-50.5	56.3-62.5	27.2-38.3	9.9-10.6
2100	36.2-44.9	51.8-57.9	38.5-46.7	10.9-11.8
2101	39.7-48.3	56.6-63.4	36.6-45.7	10.3-11.4
2102	44.6-51.3	58.3-66.2	28.5-41.4	9.8-10.6
2103	49.3-53.4	58.9-69.3	24.7-35.2	10.4-11.7
2104	36.6-45.2	52.2-60.3	37.5-44.1	10.6-11.9
Akça (C)	42.4-48.7	62.4-67.8	39.2-46.5	10.2-11.4

3633 C. Control

3634

3635 Table 3. Yield and some quality characteristics of Akça pear types.

3636 Çizelge 3. Armut tiplerinde incelenen verim ve bazı kalite özellikleri.

Type no	Cumulative yield (kg)	Cumulative yield per 1 cm ² cross-sectional area of trunk (kg /cm ²)	Average fruit weight (g)	Total soluble solids (%)
2036	27.939	0.260 bcd	90.20 a	11.0
2039	17.662	0.107 de	49.75 defg	13.0
2042	15.608	0.089 e	54.50 cde	13.2
2043	23.333	0.077 e	29.40 g	12.5
2045	60.621	0.236 abc	62.50 bcd	13.2
2051	78.912	0.350 a	73.40 abc	13.0
2099	16.532	0.087 e	51.70 def	13.5
2100	22.216	0.119 de	31.30 fg	11.0
2101	9.112	0.077 e	49.20 defg	12.5
2102	10.635	0.097 de	62.00 bcd	11.0
2103	57.724	0.338 a	81.40 ab	15.0
2104	32.115	0.136 cde	37.20 efg	11.2
Akça (C)	28.628	0.264 ab	54.00 cde	11.5

3637 C. Control

3638

3639

3640 Table 4. The ranking of Akça pear types in respect to their total points by 'weighed- ranked' method.

3641 Çizelge 4. Armut tiplerinin tartılı drecelendirmeye göre aldıkları puanlar.

Types	Yield	Fruit weight	Earliness	Total
2036	280	300	210	790
2051	400	240	120	760
2103	400	270	30	700
Akça (C)	280	150	120	550
2045	240	180	120	540
2104	120	60	300	480
2100	80	30	300	410
2039	80	120	210	410
2042	40	150	210	400
2101	40	120	210	370
2043	40	30	300	370
2099	40	120	210	370
2102	40	180	120	340

3642 C.Control



International Congress on Advances in Bioscience and Biotechnology

